

BINDING ENERGY AND THE FUNDAMENTAL PLANE OF GLOBULAR CLUSTERS

Dean E. McLaughlin¹

Department of Astronomy, 601 Campbell Hall
University of California, Berkeley, CA 94720-3411
dean@crabneb.berkeley.edu

ABSTRACT

A physical description of the fundamental plane of Galactic globular clusters is developed which explains all empirical trends and correlations in a large number of cluster observables and provides a small but complete set of truly independent constraints on theories of cluster formation and evolution in the Milky Way.

In a very good first approximation, the internal structures and dynamics of Galactic globulars are described by single-mass, isotropic King (1966) models and are thus characterized by four nominally independent physical parameters. Within this theoretical framework, it is shown that (1) 39 regular (non-core-collapsed) globulars with measured central velocity dispersions all share a common core mass-to-light ratio, $\Upsilon_{V,0} = 1.45 M_{\odot} L_{\odot}^{-1}$, and (2) 109 regular globulars both with and without direct observations of $\Upsilon_{V,0}$ show a very strong correlation between global binding energy and total luminosity, regulated by Galactocentric position: $E_b = 7.2 \times 10^{39} \text{ erg } (L/L_{\odot})^{2.05} (r_{gc}/8 \text{ kpc})^{-0.4}$. The observational scatter about either of these two constraints can be attributed fully to random measurement errors, making them the *defining equations of a fundamental plane* (FP) to which real clusters are confined in the larger, four-dimensional parameter space of general King models. They are shown to underlie a pair of bivariate correlations first found, and used to argue for the existence of a globular cluster FP, by Djorgovski (1995). A third, weaker correlation, between clusters' total luminosities and King-model concentration parameters, is related to the (non-random) distribution of globulars *on* the FP. With L , $\Upsilon_{V,0}$, E_b , and the central concentration c thus chosen as the four physical quantities that define any single globular cluster, the FP equations for $\Upsilon_{V,0}$ and $E_b(L, r_{gc})$ are used to *derive* expressions for *any other* observable in terms of only L , r_{gc} , and c . Results are obtained for generic King models and applied specifically to the globular cluster system of the Milky Way.

Subject headings: galaxies: fundamental parameters — galaxies: star clusters — globular clusters: general

¹Hubble Fellow

1. Introduction

The properties of globular clusters in the Milky Way offer empirical constraints not only on their own formation and evolution, but also on the history of our Galaxy as a whole and, to a potentially large extent, on the formation of generic star clusters—a key element of the star formation process itself. The many observed correlations between various internal structural and dynamical parameters of the clusters (core and half-light radii, surface brightnesses, velocity dispersions, etc.) that have been found by several authors (e.g., Brosche & Lentes 1984; Kormendy 1985; van den Bergh, Morbey, & Pazder 1991; van den Bergh 1994; Djorgovski & Meylan 1994; Djorgovski 1995; Bellazzini et al. 1996; Murray & Lin 1992) constitute an important set of such constraints.

The apparently large number of these correlations (see especially Djorgovski & Meylan 1994) can be somewhat misleading, however, in that it is impossible for all of them to be independent. This follows from the fact that Galactic globulars are structurally and dynamically exceedingly simple, being very well described by single-mass, isotropic King (1966) models. Since these models are defined fully by the specification of just four independent physical parameters, it is not unreasonable to expect that the same should be true, at least in a first-order approximation, of real clusters; but there can then be at most three truly distinct correlations between their various intrinsic properties (although these may be supplemented, in general, by dependences on “external” factors such as Galactocentric position or cluster metallicity).

It is clearly of interest, then, to systematically reduce the globular cluster data in the Milky Way to the minimum number of independent physical relationships required for a complete description of the observed properties of the ensemble. The establishment of such a compact set of empirical constraints—which should serve better as a target for theories of cluster formation and evolution—is the main goal of this paper. To reach it, two important results will be used as starting points.

First, the two best globular cluster correlations known to date (i.e., the only two with scatter that can be fully accounted for by observational uncertainties) are the bivariate dependences presented by Djorgovski (1995) for a subset (roughly one-third) of the Galactic globulars. One of these correlations involves only cluster core parameters, while the other includes properties measured at their half-light radii. Djorgovski found these correlations by a statistical principal-components analysis, and he offered no physical interpretation of them; but he pointed out that their very existence suggests that globular clusters may in practice be confined—much like elliptical galaxies and other large stellar systems are—to a “fundamental plane” in the higher-dimensional space of physical parameters available to them in principle.

Second, an almost completely unexamined (but, as it happens, very closely related) issue has to do with the dependence of binding energy, E_b , on mass (or luminosity) among globular clusters. This was studied some time ago by Saito (1979), who also investigated the question for dwarf and giant elliptical galaxies. He found that $E_b \propto M^{1.5}$ or so—apparently consistent with very simple

virial-theorem arguments—for globulars and giant ellipticals alike. However, his observational sample included only ten relatively bright globular clusters. The data required for a calculation of binding energy are now available for many more clusters than this and they are obviously of higher quality than those available to Saito. A total energy E_b is the only obvious physical property that is *not* routinely estimated from observations of Galactic globulars; but it is arguably important—if for no other reason than to complete a systemization of the data—to clarify the empirical behavior of this quantity.

In what follows, catalogued data are used to estimate binding energies within the context of single-mass, isotropic King (1966) models for 109 “regular” (non-core-collapsed) globular clusters in the Milky Way. The data and necessary aspects of the models are summarized in the next Section and in a pair of Appendices. Section 3 presents the binding energies themselves, and also derives core mass-to-light ratios, $\Upsilon_{V,0}$, for 39 of these clusters with measured central velocity dispersions. It is shown that (1) $\Upsilon_{V,0}$ is a constant for all clusters, and (2) E_b correlates tightly with total cluster luminosity (or mass), scaling as $E_b \propto L^{2.05 \pm 0.1}$. This is significantly (if not surprisingly) different from Saito’s (1979) result, and perhaps also from the relation obeyed by elliptical galaxies (although this also appears not to have been reviewed in any detail since Saito’s work).

Section 3 also establishes the choice of L , E_b , $\Upsilon_{V,0}$, and the King-model concentration parameter $c \equiv \log(r_t/r_0)$, as the four “independent” physical properties used to define any globular cluster. The tight constraints found on $\Upsilon_{V,0}$ and $E_b(L)$ then imply that real clusters have (at most) only two free parameters, and they do indeed occupy a fundamental plane in the theoretical space of King models. The influence of environment on this plane, as reflected in cluster Galactocentric radii and metallicities, is also examined in §3; it is shown to be confined to a dependence on r_{gc} in the *normalization* of the E_b – L scaling.

In §4, an orthogonal coordinate system called ϵ -space (analogous to the so-called κ -space of elliptical galaxies; Bender, Burstein, & Faber 1992) is constructed to obtain a face-on view of the fundamental plane. This shows that globulars are not distributed uniformly over it; essentially, the parameter c , which describes the internal density profile of a cluster, is also correlated with total luminosity or mass (see also van den Bergh 1994; Bellazzini et al. 1996). Alternate representations of the fundamental plane, in the form of a number of bivariate correlations between cluster core and half-light parameters, are also developed in §4. It is shown that the two correlations of Djorgovski (1995) are equivalent to the two constraints $\Upsilon_{V,0} = \text{constant}$ and $E_b \propto L^{2.05}$ and, thus, that Djorgovski identified the same fundamental plane that is developed here.

Section 5 and Appendix A then use the defining equations of the fundamental plane, along with manipulations based on generic properties of King model, to derive a set of simple relations that ultimately describe the variation of *any* globular cluster observable as a function of L , c , and r_{gc} *only*. These may themselves be manipulated to explain any observed correlation between any other combination of cluster variables. Thus, the fundamental plane indeed reduces a large

array of globular cluster data to a very small set of independent constraints for models of their formation and evolution. These are set out in §6, which summarizes the paper.

It should be noted that the choice of “basic” cluster properties adopted here—total luminosity (mass), binding energy, mass-to-light ratio, and concentration parameter—is not a unique possibility, nor is it necessary in any rigorous sense. It will be noted, for example, that the scaling of binding energy with luminosity is essentially equivalent to other, previously known correlations between half-light radii and luminosities ($R_h \sim L^0$; van den Bergh et al. 1991) and between core velocity dispersions and luminosities ($\sigma_0 \sim L^{0.52}$; e.g., Djorgovski & Meylan 1994). However, this does not make a description in terms of E_b any less correct or valid. In fact, this framework seems particularly amenable to theoretical studies of cluster evolution and formation both: Many simulations of the former follow the evolution of c and E_b over time, and a discussion of the $E_b(L, r_{gc})$ dependence in the latter context will be taken up in a subsequent paper (McLaughlin, in preparation).

2. Data and Model Calculations

It is well known that King’s (1966) theoretical sequence of isotropic, lowered isothermal spheres of single-mass stars (see also Binney & Tremaine 1987) provides an excellent match to the luminosity profiles of most Milky Way globulars and a good description of their internal kinematics. This specific model framework is therefore adopted for analyses throughout this paper. For self-consistency, and in order to make use of as large an empirical database as possible, the complications of velocity anisotropy and multi-mass stellar populations are not considered (although theoretical models incorporating these features have, of course, been constructed [Michie 1963; Gunn & Griffin 1979] and applied to a number of Galactic globulars [see Pryor & Meylan 1993]).

Harris (1996) maintains an on-line catalogue² of King-model parameters and other properties of 147 Galactic globular clusters; it is in part a synthesis and updating of earlier compilations by Trager, Djorgovski, & King (1993) and Djorgovski (1993a) (see also Trager, King, & Djorgovski 1995). Among other quantities, the catalogue lists the clusters’ heliocentric distances and their Galactocentric positions, r_{gc} ; their foreground reddenings, $E(B - V) = A_V/3.1$, and absolute magnitudes, M_V ; metallicities, $[Fe/H]$; projected half-light (or effective) radii, R_h ; central V -band surface brightnesses, $\mu_{V,0}$ (not corrected for extinction in the raw catalogue); scale radii,³ r_0 ,

²<http://physun.physics.mcmaster.ca/Globular.html>. The version used in this paper is the revision of 22 June 1999.

³The radii r_0 are referred to by Harris (1996), and by Trager et al. (1993) and Djorgovski (1993), as core radii, and they are denoted r_c in these catalogues. However, the tabulated values in fact correspond to the King-model scale radii r_0 defined by equation (3), and they are quantitatively different from the projected half-power radii of low-concentration clusters.

and concentration parameters, $c \equiv \log (r_t/r_0)$ (with r_t the tidal radius), derived from fits of King (1966) models to cluster surface-brightness profiles; and central luminosity densities, j_0 (in V -band $L_\odot \text{ pc}^{-3}$), obtained from the measured c , r_0 , and extinction-corrected $\mu_{V,0}$ (see, e.g., Djorgovski 1993a, and Appendix A below).

The King-model structural parameters are not available for six of the globulars in Harris’ (1996) tables; another cluster (Djorgovski 1) has no measured metallicity; and no central surface brightness is given for one more (the Pyxis cluster). These objects were therefore excluded a priori from the data set ultimately used here. Of the 139 clusters remaining, 30 are identified (following Trager et al. 1993) as having obvious or suspected post-core-collapse (PCC) morphologies, i.e., central density cusps. As Trager et al. discuss, such clusters are not particularly well described by King models—a concentration parameter of $c = 2.5$ has generally been assigned to them *arbitrarily*—and their tabulated scale radii r_0 and core surface brightnesses $\mu_{V,0}$ are most likely over- and underestimates, respectively. Thus, although most of the calculations described below have been performed for the PCC clusters using the data as given, the results should *not* be taken at face value; and while these objects are shown on many of the plots in this paper, they are never included in any more detailed analyses (e.g., least-squares fits).

This leaves 109 regular, King-model clusters as the main focus of attention. Of these, 103 have Galactocentric radii $r_{\text{gc}} < 40$ kpc; 4 have $65 \lesssim r_{\text{gc}} \lesssim 100$ kpc; and two (the faint Palomar 4 and AM1) lie at $r_{\text{gc}} \simeq 110$ –120 kpc. Estimates of the central line-of-sight velocity dispersions, $\sigma_{p,0}$, are given for roughly one-third (39/109) of this restricted sample, as well as for 17 of the 30 PCC clusters, by Pryor & Meylan (1993). These additional data are also employed here.

The recent study of Rosenberg et al. (1998) has been used to replace some of Harris’ (1996) numbers for the low-luminosity cluster Palomar 12. Specifically, from a King-model fit to star counts, Rosenberg et al. derive $c = 1.08$ and $r_0 = 0'.63$ (or 3.5 pc, for a heliocentric distance of 19.1 kpc), as compared to the $c = 1.94$ and $r_0 = 0'.20$ given by Harris. With $\mu_{V,0} = 20.6$, according to Harris, this then implies (following the procedure outlined in Appendix A below) a central luminosity density $j_0 \simeq 32 L_\odot \text{ pc}^{-3}$, lower than the previous value by a factor of roughly three. After incorporating these few changes for this one cluster, the final database is similar to that adopted by Djorgovski & Meylan (1994), and the subsets thereof in Djorgovski (1995) and Bellazzini (1998), for their recent investigations of the Milky Way globular cluster system.

Expositions of observational uncertainties can be found in Trager et al. (1993) and Pryor & Meylan (1993), and the situation may be summarized roughly as follows: scale and (projected) half-light radii are known to typical precisions of $\Delta \log r_0 \simeq \pm 0.1$ dex and $\Delta \log R_h \simeq \pm 0.1$ dex; central velocity dispersions, to within $\Delta \log \sigma_{p,0} \simeq \pm 0.07$ dex; central surface brightnesses, to $\Delta \mu_{V,0} \simeq \pm 0.3$ mag arcsec $^{-2}$; and concentrations, to $\Delta c \simeq \pm 0.2$ dex. Uncertainties in any derived quantities follow immediately from these estimates. For example, the central luminosity density is obtained from $\mu_{V,0} = \text{const.} - 2.5 \log (j_0 r_0)$ (eqs. [A2], [A3]), and thus $\Delta \log j_0 \simeq \pm 0.2$ dex. Similarly, the average surface brightness within a half-light radius is defined

as $\langle \mu_V \rangle_h = \text{const.} - 2.5 \log (L/R_h^2)$ (see eq. [A5]), so that $\Delta \langle \mu_V \rangle_h \simeq \pm 0.5 \text{ mag arcsec}^{-2}$. There are also uncertainties in cluster distance moduli— $\pm 0.15 \text{ mag}$ may be representative; see the discussion at Harris’ (1996) website—and thus in Galactocentric distances r_{gc} and total luminosities L (note that integrated apparent V magnitudes are generally good to a few hundredths of a magnitude). These are generally not dominant in the error budgets discussed here, however, and for the most part they can be ignored.

In order to refer cluster data to isotropic King models, it is necessary first to relate the observed velocity dispersion $\sigma_{p,0}$ to the model *scale velocity*, σ_0 . This latter quantity is defined through the distribution function assumed for the cluster stars, $f(E) \propto \sigma_0^{-3} [\exp(E/\sigma_0^2) - 1]$, and through the dimensionless cluster potential, $W(r) \equiv [\phi(r_t) - \phi(r)]/\sigma_0^2$ (King 1966; Binney & Tremaine 1987). $W(r)$ and a scaled density profile, ρ/ρ_0 vs. r/r_0 , follow from integrating Poisson’s equation with this distribution function and a specified value of $W_0 \equiv W(r=0)$. With r_t the radius at which $W = \rho = 0$, there is then a one-to-one relation between W_0 and the concentration parameter $c \equiv \log(r_t/r_0)$. In the limit of high concentration, the King model is a normal isothermal sphere, σ_0 is the true one-dimensional velocity dispersion inside the cluster, and $\sigma_{p,0} = \sigma_0$. For low concentrations $c \lesssim 1$, however, the outer, non-isothermal parts of a cluster influence projected quantities even at its center, and $\sigma_{p,0} < \sigma_0$. This effect is illustrated in the upper left-hand panel of Fig. 1. The numerical integrations summarized there have been used (given the observed concentrations c) to convert the central dispersions $\sigma_{p,0}$ to model σ_0 -values for the 39 regular and 17 PCC clusters in Pryor & Meylan (1993). In most cases, this is not a large correction: the Pryor & Meylan clusters all have $c > 0.75$, and $0.8 < \sigma_{p,0}/\sigma_0 < 1$. It does have the effect of adding slightly to the observational uncertainty, however: with $\Delta c = \pm 0.2$ and $\Delta \log \sigma_{p,0} = \pm 0.07$, the typical errorbar on $\log \sigma_0$ is roughly $\pm 0.09 \text{ dex}$.

The binding energies of Milky Way globulars are calculated in §3 from the basic definition (cf. King 1966; Saito 1979)

$$E_b \equiv -\frac{1}{2} \int_0^{r_t} 4\pi r^2 \rho \phi \, dr = \frac{1}{2} \int_0^{r_t} 4\pi r^2 \rho \left[\frac{GM}{r_t} + \sigma_0^2 W(r) \right] \, dr, \quad (1)$$

where M is the total mass and the sign has been chosen to make $E_b > 0$ for bound clusters. A dimensionless binding energy follows directly from this and is a function only of c (through W_0) for an isotropic and single-mass King model:

$$\mathcal{E}(c) \equiv \frac{GE_b}{\sigma_0^4 r_0} = \frac{81}{2} \frac{r_0}{r_t} \left[\int_0^{r_t} \frac{\rho}{\rho_0} \left(\frac{r}{r_0} \right)^2 \frac{dr}{r_0} \right]^2 + \frac{9}{2} \int_0^{r_t} \frac{\rho}{\rho_0} \left(\frac{r}{r_0} \right)^2 W(r) \frac{dr}{r_0}. \quad (2)$$

Here the central mass density ρ_0 is related to r_0 and σ_0 by the *model definition*

$$r_0^2 \equiv \frac{9\sigma_0^2}{4\pi G\rho_0}. \quad (3)$$

The function $\mathcal{E}$ is shown in the upper right panel of Fig. 1. Over the range $0.5 \leq c \leq 2.5$ appropriate to Galactic globulars, an observational uncertainty of $\Delta c = \pm 0.2$ translates to an r.m.s. variation of $\pm 0.2 \text{ dex}$ in $\mathcal{E}$ as well.

If the mass density profile $\rho(r)/\rho_0$ is replaced by its luminosity equivalent, $j(r)/j_0$, the light intensity I at a given projected radius R follows from the standard integral $\mathcal{I}(R) \equiv I(R)/(j_0 r_0) = 2 \int_R^{r_t} (j/j_0) (r/r_0) (r^2/r_0^2 - R^2/r_0^2)^{-1/2} dr/r_0$. The central surface density $\mathcal{I}_0$ is again a function of c only, and it is shown in the bottom left panel of Fig. 1. The full profile $\mathcal{I}(R)$ also uniquely predicts the radius $\mathcal{R}(c) \equiv R_h/r_0$ which contains half of a cluster’s total luminosity in projection. This is plotted against the concentration parameter at the bottom right of Fig. 1. Both $\mathcal{I}_0(c)$ and $\mathcal{R}(c)$ figure prominently in the analysis of §§4 and 5 below.

It will also be necessary to refer to another feature of single-mass King-model clusters, namely, the dimensionless total luminosity $\mathcal{L}(c) \equiv L/j_0 r_0^3 = \int_0^{r_t} 4\pi (j/j_0) (r/r_0)^2 dr/r_0$. This quantity is shown for a range of central concentrations in the top panel of Fig. 2. The lower half then compares the calculated luminosity— $L_{\text{mod}} = \mathcal{L}(c)j_0 r_0^3$ —to that directly observed for the 139 globular clusters of Harris (1996). The agreement for the 109 regular clusters is excellent (as was also noted by Djorgovski 1993a): $\langle \log L_{\text{mod}} - \log L \rangle = -0.017$ with an r.m.s. scatter, $\Delta = 0.25$ dex, that is less than the observational errorbar of ± 0.3 .⁴ Figure 2 can therefore be viewed as confirming that regular globular clusters are indeed well described by King models—and as showing the difficulties with PCC clusters, whose luminosities are often underestimated on the basis of their crude King-model parametrizations.

As has already been suggested, the concentration parameters measured for Galactic globular clusters span a reasonably wide range of values, from $c \simeq 0.5$ to $c \simeq 2.5$ (Harris 1996); moreover, there is a strong *systematic* quality to the variation of c from cluster to cluster (see, e.g., Fig. 6 below). Thus, Figs. 1 and 2 also serve to illustrate the *dynamical and structural non-homology* that characterizes the Milky Way globular cluster system: The relationships between global and local properties of a cluster (i.e., any of the ratios plotted in these Figures) obviously depend quantitatively on its internal density and velocity profiles. But the details of these profiles are quite sensitive to the King-model parameter c (or, equivalently, W_0), so that even *dimensionless* ratios such as $\mathcal{E}$, $\mathcal{R}$, and $\mathcal{L}$ can differ significantly between clusters; one globular cannot, in general, be turned into another simply by applying a single scaling factor to its basic physical parameters.

This is a natural consequence of the fact that globular clusters are well described by King models (it would not be the case if, for example, these objects were singular isothermal spheres), and it is clearly key to a full understanding of cluster structures and a complete explanation of the various correlations between their properties. Although this point has always been appreciated in principle (e.g., Djorgovski 1995; Bellazzini 1998), its consequences are fully explored for the first time in this paper. Throughout, the functions of c just discussed are used repeatedly to derive and manipulate physical parameters of the Milky Way globulars, thus always taking into account—or, in some sense, correcting for—any non-homology. Whenever an evaluation of $\mathcal{E}$, $\mathcal{I}_0$,

⁴This errorbar, drawn in the bottom panel of Fig. 2, follows from (conservatively) combining in quadrature the uncertainties of ± 0.1 in $\log r_0$; ± 0.12 in $\log (j_0 r_0)$; and ± 0.16 dex in $\log \mathcal{L}(c)$ (which follows from the shape of $\mathcal{L}$ in the interval $0.5 \leq c \leq 2.5$, given $\Delta c = \pm 0.2$ dex).

$\mathcal{R}$, or $\mathcal{L}$ is required, it is obtained from an observed c -value by interpolating on the numerical model curves plotted in Figs. 1 and 2. (For convenience, however, Appendix B also gives simple approximations to these dimensionless quantities as polynomial functions of the concentration c .) It is worth noting that analogous corrections for non-homology in elliptical galaxies may remove some significant part of the observed “tilt” of their fundamental plane (see Graham & Colless 1997; Busarello et al. 1997).

3. Two Basic Properties of Galactic Globular Clusters

The full spatial structure and internal dynamics of a single-mass, isotropic King (1966) model cluster are set by specifying four independent physical parameters. The four most convenient, for purposes of comparison with observations of real clusters, are the central concentration c [or the central potential depth W_0 , to fix the overall shape of a surface brightness profile $\mu(R) \sim -2.5 \log I(R)$]; the scale radius r_0 [for horizontal normalization in a plot of $\mu(R)$ vs. projected radius R]; the central luminosity j_0 (the vertical normalization, related to μ_0 as described above and in Appendix A); and the core mass-to-light ratio Υ_0 (for normalization of the model to observed line-of-sight velocity dispersions). More generally, however, *any* set of four linearly independent combinations of these quantities will serve equally well as a physical “basis” for the sequence of King models. (Such additional factors as metallicity or galactocentric position are quite separate from any model characterization of a cluster’s internal structure, and they are best viewed as “external” physical parameters.) As will be described below, the basis ultimately chosen here comprises the concentration, c ; the (logarithmic) mass-to-light ratio, $\log \Upsilon_0$; total luminosity, $\log L$ (which agrees well with the model quantity $[\log j_0 + 3 \log r_0 + \log \mathcal{L}(c)]$); and binding energy, $\log E_b = \text{const.} + [2 \log \Upsilon_0 + 2 \log j_0 + 5 \log r_0 + \log \mathcal{E}(c)]$.

At any rate, the implication is that, insofar as Galactic globulars can be approximated by the simplest King models—that is, to the extent that internal velocity anisotropies and a range of constituent stellar masses can be ignored as second-order corrections—they constitute a nominally four-parameter family of stellar systems. As was mentioned in §1, however, there exist many correlations between various properties of globular clusters in the Milky Way, implying that these objects in reality inhabit only a small part of the physical space potentially available to them in principle. Indeed, as Djorgovski (1995) and Bellazzini (1998) have suggested, and as will now be shown in a different manner, there exist at least two (and perhaps three) independent constraints linking the four basic parameters of globular clusters, so that they are confined to a fundamental plane (perhaps a line) in the theoretical 4-space of King models.

3.1. A Constant Core Mass-to-Light Ratio

As was discussed in §2, the central line-of-sight velocity dispersions $\sigma_{p,0}$ compiled for 39 regular and 17 PCC globular clusters by Pryor & Meylan (1993) have been converted to King-model scale velocities σ_0 using the concentration parameters c given by Harris (1996). With measurements of all three of σ_0 , r_0 , and j_0 thus in hand, self-consistent mass-to-light ratios $\Upsilon_{V,0}$ may be calculated for fully one-third of the Milky Way globulars via the core-fitting procedure of Richstone & Tremaine (1986): Given σ_0 and r_0 for any one cluster, equation (3) yields a core mass density ρ_0 , from which follows $\Upsilon_{V,0} \equiv \rho_0/j_0$.

Figure 3 plots ρ_0 against j_0 for the Pryor & Meylan dataset, with the regular clusters represented by circles (open symbols use the directly observed $\sigma_{p,0}$ in place of the corrected σ_0 in eq. [3]) and PCC objects shown as open squares. The straight line traces the relation $\log \rho_0 = 0.16 + \log j_0$, obtained by a least-squares fit to the non-core-collapsed clusters (filled circles) *only*. Clearly, the intercept in this equation is the mean mass-to-light ratio $\langle \log \Upsilon_{V,0} \rangle$. Moreover, the r.m.s. scatter about this fit ($\Delta = 0.2$ dex) can be attributed entirely to random observational uncertainties;⁵ there is no evidence for any significant variation in $\Upsilon_{V,0}$ from one cluster to another. (The apparent departure of the densest core-collapsed clusters from the mean line in Fig. 3 is certainly not entirely real, and it may be completely spurious: forcing a King-model fit to these objects leads to underestimates of their core densities j_0 , and thus to the underestimates of their total luminosities in Fig. 2 and to overestimates of $\Upsilon_{V,0}$ here. Once again, PCC clusters are never included in any of the quantitative analyses in this paper.)

This result has two implications. First, almost as a practical matter, it can confidently be assumed, even in the absence of direct velocity-dispersion measurements, that *all other* (non-PCC) Galactic globulars share the same mass-to-light ratio, viz.

$$\log \Upsilon_{V,0} = \log (\rho_0/j_0) = 0.16 \pm 0.03 \quad [M_\odot L_\odot^{-1}] , \quad (4)$$

where the uncertainty is just $0.2/\sqrt{39}$. This mean $\Upsilon_{V,0} = (1.45 \pm 0.1) M_\odot L_\odot^{-1}$ is the same as the direct average of the values given for individual clusters by Pryor & Meylan (1993), even though these authors define the mass-to-light ratio differently (as the ratio of central *surface* densities) and fit different (multi-mass) King models to the raw velocity and surface-brightness data. Their numbers also show no evidence for any significant dependence of $\Upsilon_{V,0}$ on j_0 , or on cluster luminosity. Figure 3 is also roughly consistent with the mean $\Upsilon_{V,0} \simeq 1.2 M_\odot L_\odot^{-1}$ found by Mandushev, Spassova, & Staneva (1991) (who, however, claim marginal evidence for a weak trend, $\Upsilon_{V,0} \propto L^{0.08 \pm 0.06}$ with large scatter).

The second consequence of Fig. 3 and equation (4) is an effective reduction of the dimensionality of the parameter space available to Galactic globular clusters, from four to three;

⁵Given $\Delta \log \sigma_0 \simeq \pm 0.09$, $\Delta \log r_0 = \pm 0.1$, and $\Delta \log (j_0 r_0) = 0.12$, and with $\Upsilon_{V,0} \propto \sigma_0^2/j_0 r_0^2$ the uncertainty $\Delta \log \Upsilon_{V,0}$ is typically ± 0.24 dex even if the constituent errors are uncorrelated.

in practice, *the core mass-to-light ratio is not a free physical variable*. This point is taken up again in §4.2 below, where it is shown that the constraint $\log \Upsilon_{V,0} \equiv 0.16$ is equivalent to one of Djorgovski’s (1995) bivariate correlations defining the globular cluster fundamental plane.

3.2. Binding Energy vs. Luminosity

Following the discussion at the beginning of this Section, the King-model parameters remaining to characterize Galactic globulars are c , r_0 , and j_0 —or any three linearly independent combinations of these quantities plus a constant $\Upsilon_{V,0}$. The remainder of this Section, as well as §4, is devoted to establishing total luminosity L (or mass $\Upsilon_{V,0}L$) and binding energy E_b as replacements for r_0 and j_0 as two of the axes in the parameter space of King models. There are several justifications for this replacement. First, there exists a previously unrecognized, but exceptionally strong, correlation between E_b and L in the Milky Way cluster system. This correlation further reduces the dimensionality of the space inhabited by real globulars, and it provides an immediate physical explanation for Djorgovski’s (1995) second equation for the fundamental plane (§4.2). Second, an emphasis on binding energy in particular facilitates comparison with the expected properties of gaseous protoclusters (McLaughlin, in preparation), i.e., this quantity is a natural focus for theoretical investigations of the origin and evolution of the globular cluster fundamental plane. And third, a view in terms of binding energy might profitably be extended to galaxies and clusters of galaxies, perhaps to yield new insight into their fundamental planes as well.

Given equations (2) and (3), E_b may be calculated for a King-model globular cluster using measurements of c and any two of σ_0 (which, again, is inferred from the directly observed $\sigma_{p,0}$ and c), r_0 , and $\rho_0 \equiv \Upsilon_{V,0}j_0$ in one of the three equivalent relations,

$$E_b(\sigma_0, r_0) = 4.639 \times 10^{45} \text{ erg} \left(\frac{\sigma_0}{\text{km s}^{-1}} \right)^4 \left(\frac{r_0}{\text{pc}} \right) \mathcal{E}(c) \quad (5a)$$

$$E_b(\sigma_0, \rho_0) = 5.995 \times 10^{46} \text{ erg} \left(\frac{\sigma_0}{\text{km s}^{-1}} \right)^5 \left(\frac{\Upsilon_{V,0}j_0}{M_\odot \text{ pc}^{-3}} \right)^{-1/2} \mathcal{E}(c) \quad (5b)$$

$$E_b(r_0, \rho_0) = 1.663 \times 10^{41} \text{ erg} \left(\frac{r_0}{\text{pc}} \right)^5 \left(\frac{\Upsilon_{V,0}j_0}{M_\odot \text{ pc}^{-3}} \right)^2 \mathcal{E}(c) . \quad (5c)$$

With $\Upsilon_{V,0}$ a constant, equation (5c) is of particular interest in that it allows for an estimate of binding energy based entirely on photometry, i.e., velocity-dispersion measurements are not explicitly required. The function $\mathcal{E}(c)$ is shown in Fig. 1 above, and it can be approximated by equation (B2) below. As was mentioned in §2, an r.m.s. variation of ± 0.2 dex in c translates to $\Delta \log \mathcal{E}(c) = \pm 0.2$; thus, the typical measurement errors in c , $\log r_0$, and $\log j_0$ result in a total observational uncertainty of ± 0.45 – 0.75 dex (depending on whether or not the individual errors are correlated) in the logarithm of E_b .

Figure 4 plots the concentrations and the binding energies, calculated using each of equations (5) (supplemented by eq. [4]), against luminosity and mass for the 39 King-model globulars (circles) and 17 PCC clusters (squares) with σ_0 known from Pryor & Meylan (1993). Note that the total masses used in this Figure are taken directly from Pryor & Meylan, who derived them from fits of anisotropic, multi-mass King models to the cluster data. These masses are therefore highly model-dependent, and in fact it is not clear what relevance they have in the analysis of this paper, which treats globulars as isotropic, single-mass clusters (in particular, the equality $M = \Upsilon_{V,0}L$ is implicit here, but it does not necessarily hold for the numbers of Pryor & Meylan 1993). Nevertheless, the right column of Fig. 4 demonstrates that correlations between cluster concentration and luminosity/mass, and between E_b and L or M , exist independently of specific model assumptions.

The top row of Fig. 4 shows a rough trend of increasing central concentration with increasing cluster luminosity. Such a correlation has also been noted by, e.g., van den Bergh (1994), Djorgovski & Meylan (1994), and Bellazzini et al. (1996), and it was recognized in essence long ago, by Shapley & Sawyer (1927). This dependence is visibly weaker than that of E_b on L , an impression that is quantified by the difference in the Spearman rank correlation coefficients s for the various relationships. (These nonparametric measures of correlation strength, specified in each panel of Fig. 4, are computed for the regular, King-model clusters [filled circles] only.) A linear dependence can be fit to these data— $c \simeq \text{const.} + 0.4 \log (L/L_\odot)$ —but the r.m.s. scatter about it is $\Delta = 0.3$, significantly larger than the observational errorbar $\Delta c = \pm 0.2$ dex. Clearly, c is not completely independent of $\log L$, but the relation between the two does not appear to be one-to-one (or, if such a strong link does exist, it is nonlinear).

On the other hand, $\log E_b$ and $\log L$ are very strongly correlated. Indeed, if the correlation coefficients s in the lower six panels of Fig. 4 are compared to those given by Djorgovski & Meylan (1994), Djorgovski (1995), and Bellazzini (1998) for a large number of empirical relations between other observables of Milky Way clusters, it can quickly be seen that *the correlation between binding energy and luminosity (or mass) is as strong as or stronger than any other correlation between properties of Galactic globulars*. In addition to this, it is clear that a simple power-law dependence of E_b on L or M completely describes the data. Least-squares fits (for the regular clusters only) of $\log E_b$ against either $\log L$ or the $\log M$ from Pryor & Meylan (1993) are drawn in Fig. 4. The r.m.s. scatter of $\log E_b$ about the straight lines in these six fits always lies within the observational errorbar (± 0.5 dex) shown in the second row of the Figure.

The three fits of $\log E_b$ against $\log L$ are statistically indistinguishable. This is only to be expected; it reconfirms the main underlying assumption here—that globular clusters are well described by isotropic, single-mass King models—and the finding of §3.1, that the core mass-to-light ratio is an essentially constant $\log \Upsilon_{V,0} = 0.16$. Figure 5 specifically compares the binding energies computed using measurements of r_0 and ρ_0 (eq. [5c]) to those calculated from σ_0 and r_0 (eq. [5a]). Since the ratio of these is just $(4\pi G \Upsilon_{V,0} j_0 r_0^2 / 9\sigma_0^2)^2$, it should be—and is—equal to unity in the mean, with an r.m.s. scatter ($\Delta = 0.4$ dex) just twice that found in Fig. 3 for the

basic identity $\Upsilon_{V,0} = \rho_0/j_0$.

If $\log E_b$ is then specified, for each of the 39 regular clusters in Pryor & Meylan’s (1993) sample, as the direct mean of the three values derived from equations (5), a least-squares regression against cluster luminosity results in

$$\log (E_b/\text{erg}) = (39.82 \pm 0.75) + (2.05 \pm 0.15) \log (L/L_\odot) , \quad (6)$$

with a Spearman rank correlation coefficient $s = 0.92$ and an r.m.s. scatter in $\log E_b$ of ± 0.43 dex. This relation is significantly different (the uncertainties in eq. [6] are $\pm 1\sigma$) from the $E_b \propto L^{1.5}$ proposed by Saito (1979).

It is perhaps worth noting that the three fits of $\log E_b$ vs. $\log M$ in Fig. 4 are also statistically identical to each other, but that they differ formally from the $E_b(L)$ fit: $E_b \propto M^{1.8 \pm 0.1}$ is indicated, as opposed to $E_b \propto L^{2.05 \pm 0.15}$. At first glance, the difference in these exponents might seem to imply a varying *global* mass-to-light ratio, $(M/L)_V \propto L^{0.14 \pm 0.11}$ or so. Such a result is obviously of very low statistical significance, however, and of dubious origin besides. It could well be completely an artifact of *a comparison between different King models*, namely, the single-mass and isotropic version used here to compute E_b , vs. the multi-mass and anisotropic variant used by Pryor & Meylan (1993) to obtain M . In addition, a more direct inspection of the Pryor & Meylan masses shows no clear evidence for any systematic variation of global $(M/L)_V$ with cluster luminosity (consistent with the lack of any such trend in $\Upsilon_{V,0}$ from Fig. 3). As a whole, then, the current data and models still show no conclusive signs of any variation in global mass-to-light ratios among Milky Way globulars. The rest of this paper will therefore discuss cluster binding energies explicitly as a function of luminosity only, and it will be taken as given that E_b scales with total mass in the same way.

Figure 6 shows the concentration c and the binding energy E_b (calculated from eq. [5c] with $\log \Upsilon_{V,0} \equiv 0.16$) as functions of luminosity for all of the 139 globulars with data taken from Harris (1996). As before, the 109 regular clusters are plotted as filled circles, and the 30 PCC objects as open squares. Evidently, the correlations seen in Fig. 4 extend to the whole of the Milky Way cluster system. In particular, the power-law nature and high statistical significance of the E_b vs. L scaling are unchanged from the smaller sample of Pryor & Meylan. It is now found that

$$\log (E_b/\text{erg}) = (39.89 \pm 0.38) + (2.05 \pm 0.08) \log (L/L_\odot) , \quad (7)$$

where the uncertainties are again $\pm 1\sigma$ estimates. The r.m.s. scatter about this fit (to the regular clusters only) is $\Delta = 0.53$ dex in $\log E_b$, consistent with the combination of purely random measurement errors in c , $\log r_0$, and $\log j_0$. With a Spearman rank statistic of $s = 0.93$, the correlation between E_b and L is better by far than those between any other cluster properties in this expanded (and now essentially complete) dataset.

The rough correspondence between c and $\log L$ seen in Fig. 4 also persists for regular, King-model clusters— $c \approx \text{const.} + 0.4 \log L$ is again indicated—but this remains less significant

(smaller s) and more scattered (the r.m.s. deviation from the straight line in the top of Fig. 6 is $\Delta \simeq 0.35$ dex, again exceeding the typical observational uncertainty in c) than the E_b – L relation.

The cluster luminosities plotted in Figs. 4 and 6 are the integrated absolute magnitudes tabulated by Harris (1996), and they are *independent* of any King-model fits; they have *not* been computed from the formula $L = \mathcal{L}(c)j_0r_0^3$. Nevertheless, Fig. 2 above demonstrated that the model luminosities do correspond very closely to the directly observed values. Thus, a point of potential concern here is that a plot of E_b against L is effectively one of $(j_0^2r_0^5)$ against $(j_0r_0^3)$, raising the possibility that any correlation might be either trivial or spurious. Figure 7 illustrates that the situation is rather more subtle than this in reality, and that the $E_b(L)$ dependence found here is indeed nontrivial. The top two panels of this Figure show $\log r_0$ and $\log(j_0r_0)$ (which again is closely related to the central surface brightness $\mu_{V,0}$) vs. total luminosity for all of Harris’ (1996) clusters. It is immediately obvious that r_0 and j_0r_0 each correlate with L —the dashed lines drawn describe $r_0 \propto L^{-0.3}$ and $j_0r_0 \propto L^{1.25}$ —but the r.m.s. scatter Δ in these plots is much larger than the observational errorbars (drawn in the upper left corners) in either case. The point of physical interest is that the *scatter* in $\log r_0$ is (anti)correlated with that in $\log(j_0r_0)$. One example of this has already been shown in Fig. 2, with its plot of $\mathcal{L}(c)j_0r_0^3$ against L ; another is now seen in the bottom panel of Fig. 7, which shows explicitly that the specific combination $(j_0^2r_0^5) \propto E_b/\mathcal{E}(c)$ is much more tightly correlated with cluster luminosity [the errorbar in the upper left, ± 0.4 dex, represents the quadrature sum of the uncertainties in $\log(j_0^2r_0^2)$ and $\log r_0^3$]. This already suggests that the relation between E_b and L is the more fundamental one, and that the (degraded) r_0 – L and j_0r_0 – L correlations are properly viewed as deriving from it.

The details of this derivation are discussed in §5 and Appendix A, along with other examples of similarly weak or scattered monovariate structural correlations in the Galactic globular cluster system. A key consideration is the effect of the clusters’ structural non-homology (see §2). Thus—for example—the bottom of Fig. 7 is still not equivalent to the lower half of Fig. 6, because $j_0^2r_0^5$ differs from E_b by a *non-constant* factor of $\mathcal{E}(c)$ (eq. [5c]). Indeed, the straight line in Fig. 7 represents the scaling $j_0^2r_0^5 \propto L^{1.7}$; it is only because $\log \mathcal{E} \sim 0.85 c$ (*very* roughly, from Fig. 1 above) and $c \sim 0.4 \log L$ that $E_b \propto L^{2.05}$ obtains in the end. Note also that the r.m.s. scatter Δ in $\log E_b$ vs. $\log L$ is actually slightly *smaller* than that in $\log(j_0^2r_0^5)$ vs. $\log L$, despite the large scatter in the c – $\log L$ correlation; this is further evidence that the $E_b(L)$ relation is nontrivial and physically significant.

There are other ways to see this dependence, however, and it might have been anticipated on the basis of some previous work. In particular, Fig. 8 shows the projected half-light radii of the clusters in Harris (1996) as a function of their total luminosities and (3D) Galactocentric radii r_{gc} . This comparison, which is not new (cf. van den Bergh et al. 1991; van den Bergh 1995), shows that R_h depends only weakly on L but tends to increase systematically with r_{gc} [the dashed line in the upper right panel of this Figure has the equation $\log(R_h/\text{pc}) = 0.23 + 0.4 \log(r_{\text{gc}}/\text{kpc})$]. Moreover, it turns out that the latter variation is responsible for what little dependence R_h does

appear to have on L : as Fig. 8 also shows (see also van den Bergh et al. 1991), the normalized quantity $R_h^* \equiv R_h(r_{\text{gc}}/8 \text{ kpc})^{-0.4}$ is essentially independent of cluster luminosity.⁶ In fact, with a rank correlation coefficient of $s \simeq -0.1$, R_h^* and L are about as close to being perfectly *uncorrelated* ($s = 0$) as E_b and L are to being perfectly correlated ($s = 1$).

The point of this is that $E_b \propto j_0^2 r_0^5 \mathcal{E}(c)$, $L = j_0 r_0^3 \mathcal{L}(c)$, and $R_h = r_0 \mathcal{R}(c)$ together give $E_b \propto (\mathcal{E}\mathcal{R}/\mathcal{L}^2) L^2/R_h$. But it happens that the non-homology factor $(\mathcal{E}\mathcal{R}/\mathcal{L}^2)$ varies very little over the range of concentrations $0.5 \leq c \leq 2.5$ appropriate for regular Galactic globulars (see Figs. 1 and 2 above, or Fig. 14 below). The simple scaling $E_b \propto L^2/R_h$, which is self-evident if the question of non-homology is ignored altogether, is then quite accurate; and the weak dependence of R_h on L —even if the influence of Galactocentric radius is also ignored—leads inevitably to the rough expectation, $E_b \propto L^2$. That is, Figs. 8 and 6 are equivalent characterizations of the Milky Way globular clusters; $E_b \sim L^2$ if and only if $R_h \sim \text{constant}$ for King-model clusters, and conceptually one constraint is no more correct or fundamental than the other. As a practical matter, however, note that the r.m.s. scatter about the mean $\log R_h^*$ in Fig. 8 is about twice the measurement uncertainty of ± 0.1 dex, whereas the scatter about the line $E_b \propto L^{2.05}$ in Fig. 6 is basically comparable to the observational errorbar. The source of the scatter in Fig. 8 is discussed in §5 (Fig. 17), which also considers another known correlation ($\sigma_0 \propto L^{0.52}$; see Fig. 18 and Djorgovski & Meylan 1994) that is equivalent to the binding energy–luminosity scaling found here.

The rest of this paper keeps its focus on binding energy, rather than half-light radius or velocity dispersion explicitly, as one of the four defining parameters of globular clusters; and the $E_b(L)$ relation, rather than the constancy of R_h , will be used (along with the constancy of $\Upsilon_{V,0}$) to define the fundamental plane. Figure 8 makes it clear that this course is not strictly necessary, but it has been adopted for the convenience and insight it offers (as has already been suggested, a discussion in terms of E_b can greatly facilitate theoretical studies of cluster formation and evolution).

Before the globular cluster fundamental plane can be constructed explicitly, however, Fig. 8 raises another issue to be addressed: Given that $E_b \propto L^2/R_h$ and $R_h \propto r_{\text{gc}}^{0.4}$, it is clear that “secondary” quantities such as Galactocentric position or cluster metallicity—which are extraneous to the formal King-model fitting process—are not completely disconnected from the basic cluster attributes $\Upsilon_{V,0}$, c , L , and E_b . What, then, is the total effect of such environmental factors on these fundamental parameters and the relationships between them?

⁶The relation claimed by Ostriker & Gnedin (1997), $R_h \propto L^{-0.63}$ for clusters with $5 \leq r_{\text{gc}} \leq 40$ kpc, is not consistent with the earlier analysis of van den Bergh et al. (1991), and it is *not confirmed* here. Again, the plot of R_h^* against L shows that *at a fixed Galactocentric radius*, cluster sizes are quite insensitive to their total masses. This point is important for discussions of cluster destruction timescales, especially in the low-mass regime.

3.3. The Influences of Galactocentric Position and Metallicity

Figure 9 shows the relative insensitivity of $\Upsilon_{V,0}$, L , and c to r_{gc} and $[\text{Fe}/\text{H}]$ for globular clusters in the Milky Way. The top row incorporates only the sample of Pryor & Meylan (1993), with $\Upsilon_{V,0} \equiv 9\sigma_0^2/(4\pi G j_0 r_0^2)$ as in §3.1. Clearly, a cluster’s core mass-to-light ratio is quite detached from its metallicity. There is perhaps a hint of a slight decrease in $\Upsilon_{V,0}$ towards larger Galactocentric radii, but this is not statistically significant. Thus, although it would be interesting to examine the question more closely by obtaining velocity data for the remaining two-thirds of the Galactic globulars, it will be assumed here that $\Upsilon_{V,0}$ is completely independent of r_{gc} . (Note that the values for the PCC clusters [open squares] are overestimates, due to the underestimation of their central densities j_0 [cf. Fig. 3]. Also, the point at $r_{gc} \simeq 90$ kpc corresponds to NGC 2419, which appears as an outlier in a plot of σ_0 against L [see Fig. 18 below], and for which $\Upsilon_{V,0}$ may have been underestimated as a result.)

The middle panels of Fig. 9 illustrate the well known facts (e.g., Djorgovski & Meylan 1994) that globular cluster luminosities are completely uncorrelated with $[\text{Fe}/\text{H}]$ —there is no mass-metallicity relation—and very weakly anticorrelated with Galactocentric position. However, it can be seen that the latter result does not reflect any clear, systematic dependence of L on r_{gc} , but appears instead to be driven by a handful of “excess” faint objects at large radii ($r_{gc} \gtrsim 10$ kpc) which are not equally represented in the inner parts of the Galaxy. This slight imbalance may be due to dynamical evolution from an initial distribution of cluster luminosities which was much more insensitive to r_{gc} (see, e.g., McLaughlin & Pudritz 1996), with a larger fraction of the low-mass clusters inside the Solar circle having been disrupted by disk shocking and evaporation over a Hubble time. Or, alternatively, since some of the outer-halo globulars are somewhat younger than average, it could be that clusters formed with lower average masses at lower redshifts (S. van den Bergh, private communication). But whatever its cause, this effect does *not* change the basic scalings inferred in Fig. 6 for concentration and binding energy as functions of luminosity.

Although a plot of c alone against r_{gc} shows a rough anticorrelation (Djorgovski & Meylan 1994), this is simply a result of the tendencies for c to decrease towards lower L ($c \approx \text{const.} + 0.4 \log L$, from Figs. 4 and 6) and for the lowest-luminosity clusters to lie preferentially at large Galactocentric radii. The bottom panels of Fig. 9 correct for this, showing that *at a given cluster luminosity*, c is essentially independent of r_{gc} . Equivalently, the mean relation between concentration and luminosity does not change with Galactocentric position (see also Figs. 12 and 13 below). It is obvious, too, that c is—like $\Upsilon_{V,0}$ and L —impervious to $[\text{Fe}/\text{H}]$.

Figure 10 shows that similar remarks apply to the *scaling*, though not to the normalization, of the $E_b(L)$ relation. The top plot does show a tendency for the lowest-energy King-model clusters to be found at the largest radii; but this is due in part to the preponderance of faint clusters in the outer parts of the Galaxy. If the overall dependence $E_b \propto L^{2.05}$ is removed, the influence of Galactocentric position alone can be isolated: the straight line in the bottom of Fig. 10 has the equation $\log(E_b/\text{erg}) - 2.05 \log(L/L_\odot) = 40.22 - 0.4 \log(r_{gc}/\text{kpc})$. The slope of this line is

uncertain by perhaps ± 0.1 ; it is, of course, just that expected from the proportionality $E_b \propto L^2/R_h$ and the systematic increase of cluster radii with r_{gc} (Fig. 8). Put another way—and as can be confirmed either by plotting $E_b r_{gc}^{0.4}$ against L or by fitting E_b vs. L directly in a series of narrow radial bins—the *slope* of the $E_b(L)$ relation in the Milky Way does not vary with Galactocentric position, but the normalization decreases systematically towards larger r_{gc} (see also Fig. 12). All the data are described completely by

$$\log (E_b/\text{erg}) = [(39.86 \pm 0.40) - 0.4 \log (r_{gc}/8 \text{ kpc})] + (2.05 \pm 0.08) \log (L/L_\odot) , \quad (8)$$

in which the errorbars are $\pm 1\sigma$ and all clusters are normalized to a common $r_{gc} = 8 \text{ kpc}$ for convenience. Note that the r.m.s. scatter about this relation, $\Delta = 0.49 \text{ dex}$, is $\sim 10\%$ lower than that in equation (7) and Fig. 6, which ignored the r_{gc} dependence. It can be attributed entirely to the effects of random measurement errors in r_0 , j_0 , and c , even under the most conservative assumption that these uncertainties are uncorrelated. Also, the intercepts in equations (6) and (7) are consistent with equation (8) given the median Galactocentric radii of the Pryor & Meylan (1993) sample of regular clusters ($\overline{r_{gc}} = 8.7 \text{ kpc}$) and the full Harris (1996) dataset ($\overline{r_{gc}} = 7.0 \text{ kpc}$).

Finally, Fig. 11 shows that metallicity plays no role in setting a cluster’s total energy, a result which is completely in keeping with the irrelevance of $[\text{Fe}/\text{H}]$ to any other aspect of the internal structure and dynamics of globular clusters (see Djorgovski & Meylan 1994 for further examples).

3.4. Summary

To a very good first approximation, Galactic globular clusters can be treated as realizations of single-mass, isotropic King (1966) models. A complete basis for the physical description of the ensemble is therefore provided by the four quantities $\log \Upsilon_{V,0}$, c , $\log L$, and $\log E_b$. While this particular choice of variables is not necessary—any set of four linearly independent King-model parameters is permissible, in principle—it is certainly sufficient. Moreover, it has the advantage of immediately revealing two strong empirical constraints on the properties of Galactic globulars (each of which will, of course, be recovered in any alternative parametrization of King-model space): (1) the core mass-to-light ratio is a constant, $\log (\Upsilon_{V,0}/M_\odot L_\odot^{-1}) = 0.16 \pm 0.03$, and (2) clusters’ binding energies are set by their total luminosities and Galactocentric positions, through $\log E_b = 39.86 - 0.4 (\log r_{gc}/8 \text{ kpc}) + 2.05 \log L$. As far as the current data can discriminate, the Milky Way cluster system appears almost *perfectly* to obey these two relations—deviations from them can be attributed entirely to random measurement errors—and it therefore populates a two-dimensional subspace in the four-dimensional volume of King models. The effects of metallicity and environment on individual cluster properties are summarized *completely* (again, within the limits of current data) by the r_{gc} -dependence in the normalization of the $E_b(L)$ relation.

Equations (4) and (8) thus *define* a fundamental plane for Galactic globulars. Various representations of this are developed in the next Section. Essentially, central concentration and total luminosity or mass are then left to determine the distribution of clusters *on* the plane; but

even this is not random, as there also exists a rough correlation between c and $\log L$ (Fig. 6). This dependence is rather weak by comparison with the constraints on $\log \Upsilon_{V,0}$ and $\log E_b$, and a one-to-one $c(\log L)$ relation has *not* yet been found in the data; but, as will also be discussed in §4, the situation does suggest the intriguing possibility that Galactic globulars may actually fall along, or at least have evolved from, something closer to a “fundamental straight line” (cf. Bellazzini 1998, who suggested something similar but with a much different physical interpretation).

4. The Fundamental Plane

For much of this Section, it will be taken as given that $\Upsilon_{V,0}$ is a constant for Galactic globulars, as in equation (4). This immediately removes the core mass-to-light ratio as a physical variable, and the parameter space remaining available to the clusters is only three-dimensional.

The preceding discussion and Fig. 6 suggest a physically transparent view of the globular cluster fundamental plane (FP) as a thin, tilted slice of $(\log L, \log E_b, c)$ space—the volume left by the constraint $\log \Upsilon_{V,0} = \text{constant}$ —which is seen edge-on along the c -axis (in the $\log L$ – $\log E_b$ plane) and closer to face-on along the E_b axis (in the c – $\log L$ plane). As was discussed in §3.3, however, the normalization of the $E_b(L)$ relation decreases systematically with increasing Galactocentric radius. Thus, a plot of $\log E_b$ vs. $\log L$ for the entire Milky Way cluster system, with its wide range of r_{gc} , actually provides an edge-on view of a *collection* of separate and distinct FPs which intersect the $\log E_b$ axis at different points. Figure 6 can therefore be improved by introducing the normalized quantity (cf. eq. [8])

$$\log E_b^* \equiv \log E_b + 0.4 \log (r_{\text{gc}}/8 \text{ kpc}) , \quad (9)$$

which simply removes all environmental influences on the FP.

Figure 12 plots c and $\log E_b^*$ against $\log L$ for the 109 regular and 30 PCC clusters catalogued by Harris (1996), divided according to whether they lie within the Solar circle ($R_0 = 8$ kpc) or outside it. Two points are apparent from this. First, it is confirmed that the r_{gc} dependence in E_b is adequately described by a scaling close to $E_b \propto r_{\text{gc}}^{-0.4}$, and that the mean increase of c with L is, although rough, essentially independent of Galactocentric position (cf. §3.3). The solid straight lines in both bottom panels follow the relation given by equation (8) above; the dashed lines show the 3σ limits on the fitted slope and intercept. Clearly, clusters at $r_{\text{gc}} < 8$ kpc and $r_{\text{gc}} > 8$ kpc obey this equation equally well. Similarly, the dashed lines in the top right panel of Fig. 12 are linear least-squares fits of c against $\log L$ (shallower slope) and of $\log L$ against c (steeper slope) for the regular clusters at $r_{\text{gc}} > 8$ kpc (excluding Palomar 1, the most obvious outlier). These same lines are then re-drawn in the top left panel of the Figure, where they are seen to be equally acceptable as crude descriptions of c vs. $\log L$ for the King-model clusters with $r_{\text{gc}} < 8$ kpc.

Second, it can be seen from Fig. 12 that *both c and $\log E_b^*$ are more strongly correlated with $\log L$ for clusters beyond the Solar circle than for those within it*: the Spearman rank correlation

coefficients are higher, and in the case of c vs. $\log L$, the r.m.s. scatter about the linear fits is smaller. This point has already been discussed in relation to the central concentrations by Bellazzini et al. (1996) and Vesperini (1997), who note that globulars at $r_{gc} < 8$ kpc have been subjected to much stronger dynamical evolution than have those at larger radii (evaporation is faster and disk shocks are more severe in denser regions of the Galaxy). Thus, the very existence of a c - $\log L$ correlation among the more distant clusters suggests that it has a largely primordial origin. These authors' quantitative numerical simulations confirm in more detail that “evolutionary processes are unlikely to play a dominant role in establishing the c - $\log M$ correlation” (Vesperini 1997). The same calculations show explicitly that the main effect of the evolution within $r_{gc} = 8$ kpc is to increase the *scatter* in the final version of an initial $c(\log L)$ dependence. This is consistent with the appearance of the data in the top half of Fig. 12. (It must be noted again, however, that the r.m.s. scatter of measured c values about the linear regressions at $r_{gc} > 8$ kpc is still substantially larger than the typical errorbar of ± 0.2 dex. It remains unclear whether there might exist a *nonlinear* c - $\log L$ correlation that is significantly tighter, or whether there is simply a real scatter in cluster concentrations at any given luminosity.)

Similarly, the very strong correlation of $\log E_b^*$ with $\log L$ for the globulars at $r_{gc} > 8$ kpc implies that this dependence, too, may have been set largely at the time of cluster formation. If this is correct, then it seems that the main effect of the stronger evolution within the Solar circle may have been a greater fractional depopulation of the low-mass tail of the cluster distribution, rather than any significant change in a power-law scaling between E_b and L . (It is emphasized again that the r.m.s. scatter in the data of both lower panels of Fig. 12 is essentially the same as, or even smaller than, the typical observational errorbar on $\log E_b$.) Indeed, an initial scaling close to $E_b \propto L^2$ can be expected to be roughly preserved during the course of dynamical evolution of an entire cluster system. Direct calculations (see the discussion and references in Murray & Lin 1992; Harris & Pudritz 1994) show that the half-mass radii of individual clusters are fairly well preserved during dynamical evolution, even over the course of a Hubble time, i.e., $dR_h/dt \approx 0$; but this and $E_b \propto L^2/R_h$ together imply $d \log E_b/dt \approx 2 d \log L/dt$. Thus, as a cluster loses mass to evaporation and tidal shocking it should move down and to the left in the $(\log L, \log E_b^*)$ plane, along a line roughly parallel to the initial correlation (if this was indeed roughly $\sim L^2$, and assuming that any changes in $0.4 \log r_{gc}$ can be neglected).

To put this another way, it is plausible that *clusters remain more or less in the fundamental plane as they evolve* and, thus, that globular clusters may have appeared on the FP as a result of the formation process. In addition, the relation between c and $\log L$ implies that the distribution of globulars over the FP is non-random and also reflects something of initial conditions. To see this properly, however, requires a further manipulation of the basic cluster variables.

4.1. ϵ -Space

By construction, the $\log L$, $\log E_b^*$, and c axes are mutually orthogonal in the three-dimensional cluster parameter space selected by $\log \Upsilon_{V,0} = \text{constant}$. And in the Milky Way, the view of the FP in a plot of $\log E_b^*$ vs. $\log L$ is nearly perfectly edge-on (again, this follows from the fact that random measurement errors can account fully for the scatter of observed clusters about the relation $\log E_b^* = 39.86 + 2.05 \log L$). The FP is therefore canted at quite a steep angle [$\arctan(2.05) = 64^\circ$] to the $(\log L, c)$ plane, and the top panels of Figs. 7 and 12 do *not* offer a truly face-on view of it; plots of c against $\log L$ suffer from strong projection effects and may not, in general, accurately reflect the true dispersion of globulars over the actual surface of the FP. (Plots of c vs. $\log E_b^*$ would be better, but that plane still makes an angle of 26° with the FP and projection effects would still be present.) In order to be as rigorous as possible—and hopefully to allow eventually for some connection with the fundamental plane(s) of other stellar systems—it is desirable to correct for this. This is easily done in a procedure analogous to that used by Bender et al. (1992; see also Burstein et al. 1997) to define a three-dimensional “ κ -space” for elliptical galaxies. The present construct for globulars is based specifically on cluster binding energies and will therefore be referred to as ϵ -space.

The easiest way to a face-on view of the FP is to define a new coordinate system by rotating the $\log L$ – $\log E_b^*$ plane through 64° (counterclockwise) about the c axis. The new (and still mutually orthogonal) axes thus obtained are proportional to

$$\begin{aligned}\epsilon_1 &\equiv \log E_b^* - 2.05 \log L \\ \epsilon_2 &\equiv 2.05 \log E_b^* + \log L \\ \epsilon_3 &\equiv c\end{aligned}\tag{10}$$

Alternatively, of course, Galactic globulars are expected, on the basis of equation (8), to lie on a line $\epsilon_1 = \text{const.}$; and the line $\epsilon_2 = \text{const.}$ is perpendicular to this in the original $\log L$ – $\log E_b^*$ plane. In this ϵ -space, then, edge-on views of the globular cluster fundamental plane are obtained by plotting ϵ_1 against either ϵ_2 or ϵ_3 , while the (ϵ_2, ϵ_3) plane provides the face-on view.

This is shown in Fig. 13 for the full sample of 139 globular clusters from Harris (1996). The two edge-on views of the fundamental plane are drawn as the bold line $\epsilon_1 = 39.86$. The scatter of the King-model clusters about this line in both the (ϵ_1, ϵ_2) and the (ϵ_1, ϵ_3) planes are just the residuals due to measurement errors. The apparent tendency for the 30 PCC clusters to fall below the FP is, again, probably spurious: the underestimation of j_0 leads to low values of E_b^* and L , and thus of ϵ_1 , for these objects; and $\epsilon_3 = c$ has been *arbitrarily* set to 2.5 for many of them.

The plot of ϵ_3 against ϵ_2 confirms that Galactic globulars are not distributed uniformly throughout their fundamental plane. The correlation of c with $\log L$ clearly carries over into one between ϵ_3 and ϵ_2 . Moreover, because the clusters do occupy a fairly narrow swath on the FP, the potential problems with projection in the c – $\log L$ correlation are actually not too severe: the dashed line drawn in Fig. 13 has the equation $\epsilon_3 = -(12.5 \pm 3.2) + (0.13 \pm 0.03) \epsilon_2$, which

corresponds (given the definition of ϵ_2 and the scaling $E_b^* \propto L^{2.05}$) to $c \simeq 0.68 \log L + \text{const.}$ (cf. Fig. 12). This particular relation follows from fitting only the regular, King-model clusters at $r_{gc} > 8$ kpc, where, again similar to the case for c vs. $\log L$, the correlation between ϵ_3 and ϵ_2 is strongest: The Spearman statistic is $s = 0.8$ for the globulars outside the Solar circle, and the r.m.s. scatter about the best linear fit is $\Delta \approx 0.3$ dex; but $s = 0.4$ and $\Delta \simeq 0.4$ dex inside 8 kpc.

Again taking the view that cluster properties in the outer halo better reflect initial conditions, it seems quite clear that Galactic globulars formed in something more like a “fundamental band” than a plane per se. Moreover, it is even plausible (but not proven) that they might have been born along a *fundamental straight line*, a one-dimensional locus in ϵ -space. (This is a term coined by Bellazzini 1998, who arrived at a similar conclusion from a very different argument. See Figs. 16 and 19 below for further discussion of Bellazzini’s interpretation of the globular cluster FP.) That is, it is conceivable that globular clusters formed through a simple process, governed almost completely by the mass and Galactocentric position of a gaseous protocluster, which naturally resulted not only in $\Upsilon_{V,0} = \text{const.}$ and $E_b^* \propto L^{2.05}$, but also in some (unknown) one-to-one relation between central concentration and luminosity.

The viability of this interpretation depends to some extent—similar to what was concluded above—on the possibility that there may be some nontrivial function of $\epsilon_3 = c$ which correlates very tightly with ϵ_2 , i.e., which shows scatter within the observational uncertainties (at least for the outer-halo clusters) and which would therefore serve better as the third axis in ϵ -space. On the other hand, while globulars at $r_{gc} > 8$ kpc should have been less influenced by dynamical evolution, it is unrealistic to expect that they could have been completely untouched by it. Thus, another possibility is that some or all of the “excess” scatter of clusters in the fundamental (ϵ_2, ϵ_3) plane is indeed irreducible, but that it arose during evolution from an initial cluster distribution that showed no intrinsic scatter. Theoretical studies of cluster evolution will be key in deciding this issue and will be of considerable importance, whatever the outcome, for theories of cluster formation.

It is obviously also of interest to understand the relation between the present ϵ -space and the corresponding κ -space defined for elliptical galaxies (Bender et al. 1992) and studied for other hot stellar systems (Burstein et al. 1997)—or, equivalently, to explain the connection between the bivariate correlations for globular clusters (Djorgovski 1995, and §4.2 just below) and the analogous relations for ellipticals (Djorgovski & Davis 1987; Dressler et al. 1987; see Pahre, de Carvalho, & Djorgovski 1998 for a comprehensive recent discussion). A potentially important difference is that one full dimension—the $\epsilon_3 \equiv c$ axis—of the observational space employed here is given over to a parameter describing the broken homology of globulars, whereas the corresponding structural factor is incorporated into all three axes of the space used for ellipticals (see Bender et al. 1992). While explicit corrections can certainly be made for the effects of non-homology in galaxies (Graham & Colless 1997; Busarello et al. 1997), these are perhaps less easily pictured in κ -space than in ϵ -space. The construction of an ϵ -space for galaxies would require the use of a single family of structural and dynamical models that is highly accurate in its description of any

individual system (in order to estimate total luminosities/masses and global binding energies by extrapolation from direct observations of the galaxy cores).

That real differences exist between the fundamental planes of globular clusters and elliptical galaxies is clear from contrasts in certain of their monovariate correlations, or projections of their FPs (e.g., Kormendy 1985; Djorgovski 1993b). What remains to be seen is the extent to which these disparities might result from fundamentally different $E_b(L)$ relations, as opposed to any number of other physical distinctions (examples being a varying core mass-to-light ratio in ellipticals [van der Marel 1991]; the presence of dark matter in the galaxies [see Kritsuk 1997]; and possibly different systematics in the structural non-homology). It may be significant, in this regard, that Busarello et al. 1997 find evidence that the *kinetic* energy per unit mass measured within an effective radius in ellipticals scales as $(E_k/M) \propto \sigma_0^{1.6}$ or so, whereas the total (E_k/M) in globular clusters can be roughly—although not perfectly—estimated as $(E_b/L) \propto L^{1.05} \sim \sigma_0^2$ (see Fig. 18 below for the relation between σ_0 and L).

4.2. Bivariate Correlations

Djorgovski (1995) used a statistical, principal-components analysis to infer the existence of two strong, bivariate correlations involving core and half-light properties of Galactic globular clusters with measured central velocity dispersions (i.e., those in the catalogue of Pryor & Meylan 1993):

$$\log \sigma_{p,0} - 0.45 \log r_0 = -(0.20 \pm 0.01) \mu_{V,0} + (4.17 \pm 0.2) \quad (11a)$$

and

$$\log \sigma_{p,0} - 0.7 \log R_h = -(0.24 \pm 0.02) \langle \mu_V \rangle_h + (4.83 \pm 0.26) , \quad (11b)$$

where $\langle \mu_V \rangle_h \equiv 26.362 - 2.5 \log (L/2\pi R_h^2)$ is the average surface brightness within a projected half-light radius; $\sigma_{p,0}$ is in units of km s^{-1} ; and r_0 , R_h are measured in pc. It was the existence of these correlations that originally led Djorgovski to conclude that the Galactic globulars are confined to a fundamental plane. But the statistical analysis does not, by itself, offer any physical insight into the FP. (Djorgovski suggested, as did Bellazzini [1998] after him, that eq. [11a] is a reflection of the “pure virial theorem” applied to cluster *cores*. However, as will be discussed further below, this is not a fully satisfactory interpretation.) Indeed, note that equations (11) appear superficially to constitute two constraints on five different observables, and thus it is not clear a priori that they need define a plane of any kind. Now, however, with a specific physical basis ($\log \Upsilon_{V,0}$, c , $\log E_b$, and $\log L$) having been chosen to describe globular clusters in a King-model framework, it is possible to reduce the number of distinct variables in these equations to just four, and to show explicitly—by comparing with the bivariate $(\sigma_0, r_0, \mu_{V,0})$ and $(\sigma_0, R_h, \langle \mu_V \rangle_h)$ correlations *expected* to arise from each of the two empirical relations defining the fundamental plane—that equation (11a) derives from equation (4) above while (11b) is a reflection of equation (6). Thus, Djorgovski’s (1995) correlations *together* identify the same globular cluster FP that has been developed here.

The basic definition of the King radius (eq. [3]), together with a constant $\log \Upsilon_{V,0} = 0.16$ and the definition of surface brightness, yields a relation between σ_0 (*not* exactly the $\sigma_{p,0}$ used by Djorgovski in establishing his correlations), r_0 , and $\mu_{V,0}$. The details of this derivation are given in Appendix A; from equation (A9),

$$\log \sigma_0 - 0.5 \log r_0 = -0.2 \mu_{V,0} - 0.5 \log \mathcal{I}_0(c) + 4.241 . \quad (12a)$$

Here and throughout, σ_0 is in units of km s^{-1} , r_0 is in pc, and $\mu_{V,0}$ is in mag arcsec^{-2} . The function $\mathcal{I}_0$ is the dimensionless central surface density for a King model of concentration c ; it is given in Fig. 1 above and in equation (B3) below. An equivalent expression, in terms of cluster half-light radii and surface brightnesses, follows from defining $\langle \mu_V \rangle_h$ and from accounting for non-homology in the ratio $\mathcal{R}(c) \equiv R_h/r_0$ (see Fig. 1 and eq. [B4]); according to equation (A11),

$$\log \sigma_0 - 0.5 \log R_h = -0.2 \langle \mu_V \rangle_h - 0.5 \log [\mathcal{L}(c)/\mathcal{R}(c)] + 4.640 , \quad (12b)$$

where $\mathcal{L}$ is the dimensionless luminosity given in Fig. 2 and equation (B5).

Similarly, the analysis in Appendix A (see eqs. [A12] and [A13]) shows that the finding $\log (E_b/\text{erg}) = 39.82 + 2.05 \log (L/L_\odot)$ (eq. [6]; for clusters in the catalogue of Pryor & Meylan 1993 specifically) is equivalent to

$$\log \sigma_0 - 0.775 \log r_0 = -0.205 \mu_{V,0} + 0.25 \log \left[\mathcal{L}(c)^{2.05} / \mathcal{E}(c) \mathcal{I}_0(c)^{2.05} \right] + 3.943 \quad (13a)$$

(with the dimensionless energy $\mathcal{E}$ again discussed in §2 and Appendix B), and to

$$\log \sigma_0 - 0.775 \log R_h = -0.205 \langle \mu_V \rangle_h - 0.25 \log [\mathcal{E}(c)/\mathcal{R}(c)] + 4.352 . \quad (13b)$$

Note that, in general, the r_{gc} dependence in the normalization of the $E_b(L)$ relation (eq. [8]) should appear as an additional term ($-0.1 \log r_{\text{gc}} + \text{constant}$) on the right-hand side of equations (13). In order to make a more direct comparison with the analysis of Djorgovski (1995), however, the effect (which is evidently quite small anyway) has been acknowledged only implicitly here, by using a normalization of the E_b – L scaling that is appropriate for the median Galactocentric position of the globulars in the Pryor & Meylan database.

Figure 14 confirms the validity of equations (12b) and (13b) as a representation of the globular cluster FP in terms of half-light quantities. The left panels here show the non-homology functions $0.5 \log (\mathcal{L}/\mathcal{R})$ and $0.25 \log (\mathcal{E}/\mathcal{R})$, as obtained in general from numerical integrations of King models (curves) and evaluated specifically for the central concentrations of the 39 regular clusters (points) in the sample of Pryor & Meylan (1993). The functions vary only slightly among these clusters, and the effects of non-homology happen to be effectively suppressed in this projection of the FP. The mean values indicated for each function may therefore be applied to derive nearly constant intercepts in equation (12b) for $\Upsilon_{V,0} = \text{const.}$ and equation (13b) for $E_b \propto L^{2.05}$. The right panels of Fig. 14 then make a direct comparison between the resulting FP “predictions” (solid lines) and the Pryor & Meylan data. The agreement is clearly quite good and, as expected,

the r.m.s. scatter of the data about the expected correlations is within the range of observational uncertainty.

The top panels of Fig. 14, and equation (12b), also represent the basic scaling $L \propto (\sigma_0^2 R_h)(\mathcal{L}/\mathcal{R})$, which ultimately—given that $\Upsilon_{V,0}$ is a constant independent of L —is just an expression for the total *mass* of a King model. Since the ratio $\mathcal{L}/\mathcal{R}$ does not vary particularly strongly or systematically with c (or, therefore, with L), this result appears to differ significantly from the $L \propto (\sigma_0^2 R_h)^{0.7-0.8}$ that Schaeffer et al. (1993) find for elliptical galaxies and Abell clusters (although note that these authors claim that their scaling holds for Galactic globulars as well). This is another indication that the globular cluster fundamental plane may indeed differ from those of these other systems. Whether this particular contrast is due to fundamentally different trends of $\Upsilon_{V,0}$ with L , or to different structural systematics (i.e., to the non-homology term $\mathcal{L}/\mathcal{R}$ perhaps being more sensitive to L in the larger systems), is unclear at this point.

The two left panels of Fig. 14 further show the product $\mathcal{E}\mathcal{R}/\mathcal{L}^2$ to be very nearly constant as a function of c , and thus they confirm that the expectation $E_b \propto L^2/R_h$ for homologous clusters actually is a good approximation for King-model globulars as well (as was discussed in §3.2 above; see Fig. 8).

Figure 15 illustrates the accuracy of equations (12a) and (13a) for the globular cluster FP. Again, the panels on the left show the appropriate non-homology functions for generic King models and for the regular clusters of Pryor & Meylan (1993). The top plot shows (as does Fig. 1) that $\mathcal{I}_0$ is a very weak function of c . Non-homology is therefore a rather minor concern for this FP equation as well, and the mean value of $\mathcal{I}_0$ can be used to estimate a roughly constant intercept for equation (12a); this then compares very favorably to the (non-PCC) data in the top right panel of Fig. 15. The bottom panels, however, show that things are not always so simple. The logarithm of $\mathcal{L}^{2.05}/\mathcal{E}\mathcal{I}_0^{2.05}$ in the left panel increases systematically (and close to linearly, as the dashed line indicates) with concentration parameter c , and hence with cluster luminosity (and, thus, with central surface brightness as well; see eqs. [14] below). This function is therefore *not* well approximated by a constant, and if it is treated as such—if it is simply assigned its mean value in equation (13a), as was done in the other examples here—the neglected non-homology manifests itself as a slope in the observed correlation between $(\log \sigma_0 - 0.775 \log r_0)$ and $\mu_{V,0}$ which is steeper than the naive expectation.

Equation (13b) for $(\log \sigma_0 - 0.775 \log R_h)$ vs. $\langle \mu_V \rangle_h$ —a result of $E_b \propto L^{2.05}$ —comes closest to the basic form of Djorgovski’s (1995) equation (11b). The fact that his analysis zeroed in on a half-light correlation with the slightly different combination $(\log \sigma_{p,0} - 0.7 \log R_h)$ may itself be understood as a consequence of the fact that $E_b \sim L^{2.05}$ for the Galactic globulars: since R_h is then so nearly constant as a function of luminosity, a correlation between $0.7 \log R_h$ and $\langle \mu_V \rangle_h$ is not significantly worse, statistically, than one involving $0.775 \log R_h$. Meanwhile, equation (12a) for $(\log \sigma_0 - 0.5 \log r_0)$ vs. $\mu_{V,0}$ is clearly equivalent to equation (11a) from Djorgovski (1995),

which therefore is another statement of the fact that the King-model clusters have an essentially constant $\Upsilon_{V,0} = (1.45 \pm 0.10) M_{\odot} L_{\odot}^{-1}$.

Finally, as was noted above, it has been claimed (e.g., Djorgovski 1995; Bellazzini 1998; see also Djorgovski & Meylan 1994) that equation (11a) (or eq. [12a]) might follow from globular cluster *cores*, viewed as dynamically distinct entities in their own right, being “virialized” with a constant mass-to-light ratio. However, this is not entirely accurate. If cores are defined as those parts of clusters within the volume $r \leq r_0$, direct integrations of King models show that they do not satisfy the simplest version of the virial theorem; that is, if kinetic energy is denoted E_k , then $2E_k(r_0) \neq E_b(r_0)$ in general. This is shown in Fig. 16, which plots the ratio $2E_k/E_b$ as a function of r/r_0 for three representative King models. [Binding energy is defined, as in eq. (1), by $E_b(r) = -(1/2) \int_0^r 4\pi r^2 \rho \phi dr$; the kinetic energy is computed from $E_k(r) = (3/2) \int_0^r 4\pi r^2 \rho \sigma^2 dr$.] This may explain why Bellazzini (1998), who assumed $2E_k/E_b \equiv 1$ at $r = r_0$, inferred a mass-to-light ratio of only $\Upsilon_{V,0} \simeq 0.7 M_{\odot} L_{\odot}^{-1}$ —a factor of two too small—from his analysis of Djorgovski’s (1995) bivariate correlation for cluster cores. This Section has shown that, while $\Upsilon_{V,0}$ is indeed a constant in Galactic globulars, the precise form of the correlation between $(\log \sigma_0 - 0.5 \log r_0)$ and $\mu_{V,0}$ depends on the definition of r_0 in equation (3); but although the basic scaling there does arise generically from a dimensional analysis of Poisson’s equation (or, equivalently, from the virial theorem *including surface terms* which vanish only at the *tidal* radius), the normalization (which helps set the intercept in eq. [11a] or eq. [12a]) is a *convenience* specific to King (1966) models.

5. Other Correlations

Sections 3 and 4 have presented the main results of this paper: The binding energies of 109 regular globular clusters in the Milky Way, calculated within the theoretical context of single-mass, isotropic King (1966) models, correlate very tightly with total luminosities, and decrease systematically with increasing Galactocentric radius (eq. [8]). This result and the fact (eq. [4]) that the core mass-to-light ratio is a constant (at least for the 39 regular clusters where it has been directly measured) then imply the existence of a fundamental plane for globular clusters, one which has an immediate and clear interpretation even while accounting for the bivariate cluster correlations discovered by Djorgovski (1995). This new physical view of the FP is expected to aid in developing theories of cluster formation and evolution.

Meanwhile, on perhaps a more pragmatic note, it is a mathematical necessity, requiring no further proof once a King-model framework has been adopted, that *any correlation* between *any set of cluster observables* can be obtained simply by treating only r_{gc} , L and (to rather a lesser degree) c as independent variables and then manipulating the constraints on $E_b(L, r_{gc})$ and $\Upsilon_{V,0}$ according to generic properties of King models. (As was concluded in §3.3, all cluster attributes are independent of metallicity, a fact which—although interesting in its own right—justifies the neglect of $[\text{Fe}/\text{H}]$ in this discussion.) No other empirical trend contains any physical information

beyond the intrinsic properties of these models and the defining equations (4) and (8) of the globular cluster FP. This point is developed quite generally in Appendix A, and it is now illustrated in brief for the Milky Way cluster system specifically.

Equations (A14)–(A19) give expressions for various physical quantities in terms of the basis chosen in this paper for King-model parameter space, i.e., as functions of L , c , $\Upsilon_{V,0}$, and E_b , allowing for arbitrary $\Upsilon_{V,0}$ and any power-law scaling $E_b = A(L/L_\odot)^\gamma$. Applied to the Milky Way cluster system, with $\log \Upsilon_{V,0} \equiv 0.16$, $\gamma = 2.05$, and $\log(A/\text{erg}) = 39.86 - 0.4 \log(r_{\text{gc}}/8 \text{ kpc})$, the analysis of Appendix A therefore results in the following observable dependences on L , r_{gc} , and c :

$$\log r_0 = -0.05 \log L + 0.4 \log(r_{\text{gc}}/8 \text{ kpc}) - \log(\mathcal{L}^2/\mathcal{E}) + 1.681 \quad (14a)$$

$$\log R_h = -0.05 \log L + 0.4 \log(r_{\text{gc}}/8 \text{ kpc}) - \log(\mathcal{L}^2/\mathcal{E}\mathcal{R}) + 1.681 \quad (14b)$$

$$\log j_0 = 1.15 \log L - 1.2 \log(r_{\text{gc}}/8 \text{ kpc}) + \log(\mathcal{L}^5/\mathcal{E}^3) - 5.042 \quad (14c)$$

$$\mu_{V,0} = -2.75 \log L + 2 \log(r_{\text{gc}}/8 \text{ kpc}) - 2.5 \log(\mathcal{L}^3\mathcal{I}_0/\mathcal{E}^2) + 34.766 \quad (14d)$$

$$\langle \mu_V \rangle_h = -2.75 \log L + 2 \log(r_{\text{gc}}/8 \text{ kpc}) - 5 \log(\mathcal{L}^2/\mathcal{E}\mathcal{R}) + 36.761 \quad (14e)$$

$$\log \sigma_0 = 0.525 \log L - 0.2 \log(r_{\text{gc}}/8 \text{ kpc}) - 0.5 \log(\mathcal{E}/\mathcal{L}) - 1.872 \quad (14f)$$

for cluster radii in pc, luminosities in $L_\odot$, surface brightnesses in mag arcsec^{-2} , luminosity densities in $L_\odot \text{ pc}^{-3}$, and velocities in km s^{-1} .

Djorgovski & Meylan (1994) present many “monovariate” correlations for Galactic globulars, of the type $\log r_0$ vs. $M_V = 4.83 - 2.5 \log L$; $\log \sigma_0$ vs. $\mu_{V,0}$; $\log j_0$ vs. $\log r_{\text{gc}}$; and so on. All of these can be traced back to some combination of equations (14), and thus to King-model definitions and the two fundamental-plane relations. (Additional quantities considered by Djorgovski & Meylan include dynamical relaxation times, which are derived from measurements of other cluster parameters and thus could easily be included in the analysis here.) Clearly, however, the separation of L - and r_{gc} -dependences in many cases might give a skewed or degraded impression of the true, underlying physical relationship responsible for a correlation; and so can the neglect of structural and dynamical non-homology, i.e., of the increase of (and scatter in) c as a function of L , and the concomitant variations in $\mathcal{L}(c)$, $\mathcal{E}(c)$, $\mathcal{R}(c)$, and $\mathcal{I}_0(c)$ from one cluster to another.

One example of this is shown in Fig. 17, which follows up the FP prediction of equation (14a) to improve significantly on the rather poor anticorrelation between scale radius and luminosity that was seen in Fig. 7 above for the full set of clusters taken from Harris (1996). It is now clear that the only reason *any* such correlation appears is because the non-homology term $\log[\mathcal{L}(c)^2/\mathcal{E}(c)]$ increases (very roughly) as $\text{const.} + 0.7c \approx \text{const.} + 0.3 \log L$ (top panel of Fig. 17), far outweighing the explicit dependence on $\log L$ in equation (14a). The correlation plotted in the bottom panel of this Figure takes this and the expected dependence on Galactocentric position fully into account. The solid line drawn through the data here is just equation (14a). The dashed line is a formal least-squares fit to the 109 regular-cluster points. Although it has the expected slope of -0.05 , its

intercept is offset slightly downwards, a discrepancy which traces back to the small—and purely observational—bias introduced by assuming that $L = \mathcal{L}(c)j_0r_0^3$ exactly for the Galactic globulars (see the discussion around Fig. 2). The r.m.s. scatter of the data points about either of these lines is a factor of almost 2 smaller than in a plot of $\log r_0$ against $\log L$ alone and appears to be fully attributable to random measurement errors.⁷

It is worth noting that this approach also explains the “large” scatter of $\log R_h^*$ values in Fig. 8 above ($\Delta = 0.21$ dex, as opposed to the observational errorbar of ± 0.1 on $\log R_h$ alone): the scatter about the prediction of equation (14b) is guaranteed to be the same as that about equation (14a), since the former comes from adding $\log \mathcal{R}(c)$ to each side of the latter.

All of the other equations above can be similarly dissected, but it is particularly worth looking at the correlation of velocity scale with total luminosity (eq. [14f]). This is compared to cluster data from the smaller sample of Pryor & Meylan (1993) in Fig. 18. In the top panel, non-homology and the expected variation with Galactocentric radius are ignored. A fairly strong correlation persists in this case, however, because the ratio $\mathcal{E}/\mathcal{L}$ depends only weakly on c (as can be inferred from the similar shapes of the individual curves in Figs. 1 and 2 above) and because all but one of the regular (non-PCC) Pryor & Meylan clusters lie within $2.6 \lesssim r_{\text{gc}} \lesssim 29$ kpc [so that the term $0.2 \log(r_{\text{gc}}/8 \text{ kpc})$ ranges only from -0.1 to $+0.1$ or so]. The exception to this is the bright cluster NGC 2419, marked on Fig. 18, which is located at $r_{\text{gc}} = 91.5$ kpc according to Harris (1996). This partly explains its appearance as an outlier relative to the straight line drawn through the top panel of the Figure:

$$\log \sigma_0 = 0.525 \log L - 1.928 ,$$

which follows from applying an average $0.5 \log(\mathcal{E}/\mathcal{L}) \simeq 0.049$ and a median $r_{\text{gc}} = 8.7$ kpc (both evaluated using the 39 regular clusters in the sample) to equation (14f). Clearly, this simple version of the full relation already provides a reasonable description of the data (aside from NGC 2419), and the monovariate correlation is a rather faithful representation of the FP scaling $E_b^* \propto L^{2.05}$ and the constancy of $\Upsilon_{V,0}$.

The situation nevertheless improves somewhat if $\mathcal{E}/\mathcal{L}$ and r_{gc} are treated properly as variables. This is done in the bottom panel of Fig. 18, which shows a stronger and more significant σ_0 – L correlation (i.e., the r.m.s. scatter Δ is lower and the Spearman rank correlation coefficient s is higher), and in which NGC 2419 appears as a less extreme outlier (though an outlier all the same). The solid line plotted here traces the full equation (14f), and it is statistically indistinguishable from a formal least-squares fit to the regular clusters (dashed line). The scatter of these data

⁷The observational errorbar on $\log r_0$ is ± 0.1 dex, and an uncertainty of ± 0.2 in c translates to about ± 0.15 in $\log(\mathcal{L}^2/\mathcal{E})$. In addition, the derivation of equation (14a) in Appendix A involves squaring the equality $L = \mathcal{L}j_0r_0^3$, thereby introducing additional scatter of $2 \times 0.25/\sqrt{109} \simeq 0.05$ dex about the mean line (see Fig. 2). Even added conservatively, in quadrature, these three terms lead to an expected r.m.s. scatter of $\Delta \simeq 0.2$ dex, essentially that found in the bottom of Fig. 17.

about either line is again that expected given the typical observational uncertainties.⁸

As was mentioned above, equations (14) can be combined in various ways to produce a large number of other correlations. For instance, the sum of (14a) and (14c) gives the full dependence underlying the rough correlation between $\log(j_0 r_0)$ and $\log L$ seen in Fig. 7 above. Another example—and the last to be considered here—is found by eliminating $\log r_{gc}$ between equations (14a) and (14d) above:

$$\log r_0 - 0.2 \mu_{V,0} = 0.5 \log L - 0.5 \log (\mathcal{L}/\mathcal{I}_0) - 5.272 . \quad (15)$$

Alternatively, the manipulations leading to equation (A16) below show that this relation is really just an expression for the total luminosity of a King model, combined with the definition of surface brightness; as such, it is completely independent of any fundamental plane specifications.

Equation (15) relates to the correlation between scale radius and core surface brightness, shown in the top left panel of Fig. 19, for the 109 regular and 30 PCC clusters from the catalogue of Harris (1996). This is of interest because Bellazzini (1998) has claimed that the correlation (along with one between σ_0 and $\mu_{V,0}$, which can actually be obtained from eqs. [14d] and [14f]) results from Galactic globulars having a “constant core mass.” However, this conclusion was based on the assumption that cores satisfy the virial theorem in the form $2E_k(r_0) = E_b(r_0)$, and this was shown above (Fig. 16) *not* to be true in general. Moreover, the basic hypothesis of a constant $M_0 \equiv \Upsilon_{V,0} j_0 r_0^3$ is not borne out by the data. The top right panel of Fig. 19, for instance, shows clearly that measured values of $j_0 r_0^3$ range over more than two orders of magnitude—far in excess of the observational errorbar—in the Milky Way cluster system. Bellazzini (1998) also recognizes this but implies, in essence, that the scatter, although unarguably real, is consistent with more or less random excursions from a line of constant core mass. However, the bottom left panel of Fig. 19 shows that this is not the case either: If it were true that $j_0 r_0^3 \sim \text{const.}$, such that $r_0 \sim (j_0 r_0)^{-1/2}$ with some intrinsic scatter, then the quantity $(\log r_0 - 0.2 \mu_{V,0})$ would have to be *uncorrelated* with any other cluster property; but instead it shows a significant dependence on the total cluster luminosity.

The point is that *both* the monovariate r_0 – $\mu_{V,0}$ correlation, *and* a correlation between $\log r_0 - 0.2 \mu_{V,0}$ and $\log L$, are expected on the basis of equation (15). But this says nothing about a constant cluster core mass; rather, it implies that King-model clusters at a given *total* luminosity (and with a single central concentration) will always show a correlation of the form $\log r_0 \sim 0.2 \mu_{V,0}$, *by definition*. The rougher trends and the large scatter in the top half of Fig. 19 reflect the superposition of many such correlations for clusters with a wide range of total L in the Milky Way, modified somewhat by systematics in the non-homology terms $\mathcal{L}$ and $\mathcal{I}_0$ that result from the range in c -values as well.

⁸In this case, the errorbar on $\log \sigma_0$ is ± 0.09 dex (§2). However, the intercept in equation (14f) includes a contribution from $0.5 \log \Upsilon_{V,0}$ (see the general eq. [A19]), which is further subject to errors in $\log \sigma_0$ (because $\Upsilon_{V,0} \propto \sigma_0^2$; see §3.1). The total expected scatter in Fig. 18 is therefore $2 \times \Delta(\log \sigma_0) = 0.18$.

The solid line in the bottom left of Fig. 19 is just equation (15) with an intercept evaluated assuming a constant $\log(\mathcal{L}/\mathcal{I}_0) = 1.007$, the average for Harris’ (1996) King-model clusters; the dashed line is a least-squares fit to the data. The two differ—and the scatter about either exceeds the observational errorbars on the data—because the ratio $\mathcal{L}/\mathcal{I}_0$ has a significant dependence on c , and hence on $\log L$ (see Figs. 1 and 2); the cluster non-homology cannot be ignored here. The bottom right corner of Fig. 19 therefore shows the full correlation expected on the basis of equation (15). The agreement with the data is now excellent; the solid, “model” line is indistinguishable from a least-squares fit to the regular clusters, and the r.m.s. scatter is fully within the realm of random measurement errors. [This last plot is equivalent to one of $(\mathcal{L}j_0r_0^3)^{1/2}$ against $(L/L_\odot)^{1/2}$, so the observed and expected scatter both are just half those in the bottom panel of Fig. 2 above.]

Thus, the observed correlation between r_0 and $\mu_{V,0}$ in globular clusters is controlled entirely by the generic behavior of total luminosity in King models, and by the systematic (but scattered) increase of c with L found in the Milky Way. There is no need to postulate a constant core mass. Indeed, much more generally, in the scheme developed here—where globulars are defined completely by L , c , $\Upsilon_{V,0}$, and E_b^* —there is no *room* for such new constraints not already provided by the equations of the fundamental plane.

6. Summary

If they are described by single-mass, isotropic King (1966) models, globular clusters are fully defined, in general, by specifying just four independent physical parameters. These were chosen in this paper to be a mass-to-light ratio ($\log \Upsilon_{V,0}$), total binding energy ($\log E_b$), central concentration [$c = \log(r_t/r_0)$], and total luminosity ($\log L$). It has been shown that (1) all 39 regular (non-core-collapsed) Galactic globular clusters with measured core velocity dispersions have a common mass-to-light ratio (§3.1),

$$\log(\Upsilon_{V,0}/M_\odot L_\odot^{-1}) = 0.16 \pm 0.03 ;$$

and (2) if it is assumed that this also holds for the other regular clusters in the Milky Way with no velocity data, but for which measured King-model structural parameters are available, the full cluster system (109 objects in all, excluding core-collapsed members) shows a very well-defined dependence of binding energy on total luminosity, modulated by Galactocentric radius (§3.2):

$$\log(E_b/\text{erg}) = [(39.86 \pm 0.40) - 0.4 \log(r_{gc}/8 \text{ kpc})] + (2.05 \pm 0.08) \log(L/L_\odot) .$$

The scatter about each of these relations is fully accounted for by the typical observational errorbars on $\log \Upsilon_{V,0}$ ($\simeq 0.2$ dex) and $\log E_b$ ($\simeq 0.5$ dex).

With $\Upsilon_{V,0}$ essentially a fixed constant and E_b known precisely (at a given Galactocentric position) as a function of L , only two of the four basic cluster properties are truly independent; globular clusters in the Milky Way are confined to a narrow, two-dimensional subregion—a

fundamental plane (FP)—in the larger, four-dimensional space of King models. The distribution of clusters on the plane is then determined essentially by their c - and L -values. They are not scattered randomly, however, as there also exists a correlation between c and $\log L$. Still, this does *not* necessarily mean that the locus of globular clusters is only one-dimensional: a linear c - $\log L$ relation is of poorer quality than either of the empirical constraints on $\log \Upsilon_{V,0}$ and $\log E_b$, and although there *might* exist some more complicated function which gives a comparably one-to-one dependence of concentration on luminosity, one has yet to be found (§§3 and 4).

Regardless, the mean trend of c vs. $\log L$ is independent of Galactocentric position and cluster metallicity, as is the value of $\log \Upsilon_{V,0}$ and the slope of the correlation between $\log E_b$ and $\log L$ (§3.3). The normalization of the $E_b(L)$ scaling is also independent of $[\text{Fe}/\text{H}]$, but it decreases towards larger Galactocentric radii. The equation above accounts for this, and the r_{gc} dependence there describes the *full extent of environmental influences* on the globular cluster FP.

It was shown in §4 that cluster concentration parameters (and, to a lesser extent, binding energies) correlate more tightly with luminosity for globulars at $r_{\text{gc}} > 8$ kpc than for those at smaller radii. Thus, since dynamical evolution has likely been less effective at erasing initial conditions outside the Solar circle—and since simple arguments indicate that clusters could remain on or close to the FP as they evolve—it may well be that most properties of the fundamental plane were set more or less at the time of globular cluster formation. Quantitative calculations of cluster evolution over a Hubble time in the Galactic potential should be applied to check this.

A three-dimensional “ ϵ -space” was constructed in §4.1 from a simple transformation of the cluster parameter space left after specifying $\log \Upsilon_{V,0} = \text{constant}$ (and after removing the effects of Galactocentric radius) in order to obtain a directly face-on view of the FP. It could be interesting to attempt a similar construction for elliptical galaxies and clusters of galaxies, i.e., to interpret their fundamental planes in terms of binding energy as well, in order to make a more direct comparison with the globular cluster FP.

The equations for $\Upsilon_{V,0}$ and for E_b as a function of L , which define the FP, were shown in §4.2 to be equivalent to the two strong bivariate cluster correlations used by Djorgovski (1995) to argue for the existence of a fundamental plane for globular clusters. Djorgovski (and Bellazzini 1998) offered a simple virial-theorem argument (which was shown here to be incomplete) as an explanation for one of these correlations, but did not interpret the other. The results of this paper have put his results on a firm physical footing.

Finally, since $\Upsilon_{V,0}$, c , E_b , and L completely define globular clusters, any correlations connecting any of their other properties, or any other trends with Galactocentric radius, necessarily derive from generic properties of King (1966) models combined with the empirical FP relations and their environmental dependence. Appendix A outlined such derivations quite generally, and specialized results for the Milky Way cluster ensemble were presented in §5.

The picture that has been developed here is a simplification in that it works specifically in the context of single-mass and isotropic models for globular clusters. This approximation is evidently

an excellent one, but because of it the results of this paper can say nothing about the stellar mass functions or possible velocity anisotropy in observed clusters. Nor have the kinematics of clusters within the Milky Way (e.g., orbital eccentricities, or bulk rotation of the metal-rich subsystem) been considered in any way. These issues aside, however, the preceding discussion may be distilled into a small set of four main facts to be explained by theories of cluster formation and evolution:

- All cluster properties are independent of metallicity.
- The core mass-to-light ratio, $\Upsilon_{V,0} = (1.45 \pm 0.1) M_{\odot} L_{\odot}$, does not vary significantly with cluster luminosity or Galactocentric position.
- The concentration parameter, which controls the shape of a cluster’s internal density profile, correlates with luminosity. The dependence of c on $\log L$ is much more significant outside the Solar circle than inside, but even there the relationship is not obviously one-to-one. Its basic form is, however, independent of r_{gc} .
- Binding energy is intrinsically a function of luminosity and is regulated by Galactocentric position: $E_b \propto L^{2.05} r_{gc}^{-0.4}$. Since there is no evidence for significant variations in the *global* mass-to-light ratios of clusters in the Milky Way, E_b is inferred to scale with total mass in the same way as with luminosity.

Once again, the balance of current evidence suggests that these characteristics of the fundamental plane—and, thus, most of the systematics of the Galactic globular cluster system—were essentially fixed by the cluster formation process. Moreover, it is not implausible that this process was controlled largely by a single intrinsic protocluster parameter—the initial gas mass—and adjusted by an external influence depending on Galactocentric radius. Further discussion along these lines is left for future work (McLaughlin, in preparation). Regardless of any interpretive details, however, it clearly will be important to determine the extent to which the globular cluster systems of other galaxies can be described in the same simple terms that apply in the Milky Way.

This work was supported by NASA through grant number HF-1097.01-97A awarded by the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc., for NASA under contract NAS5-26555.

A. KING MODELS AND GLOBULAR CLUSTER CORRELATIONS

The analysis in this paper refers to eleven physical parameters which may be observed or derived for globular clusters: the central line-of-sight velocity dispersion $\sigma_{p,0}$; the scale radius r_0 , which is closely related (but not always identical) to the core radius, at which the luminosity

surface density falls to half its central value; the central surface brightness $\mu_{V,0}$, which is related to the central V -band intensity, I_0 , in units of $L_\odot \text{ pc}^{-2}$; the central luminosity density j_0 ($L_\odot \text{ pc}^{-3}$); the central mass-to-light ratio $\Upsilon_{V,0}$, which gives the central *mass* density, $\rho_0 = \Upsilon_{V,0} j_0$, and which must equal the global mass-to-light ratio in (idealized) clusters of single-mass stars; the projected half-light radius R_h and the surface brightness $\langle \mu_V \rangle_h$ averaged within that aperture; the central concentration c , which describes the global shape of the cluster’s surface brightness profile; the total luminosity L ; and the cluster binding energy, E_b .

There are, of course, other quantities of interest (e.g., core and half-mass relaxation times; average velocity dispersion within the half-light radius), but within the context of King’s (1966) model for globular clusters, these other parameters can all be derived from some subset of the eleven just listed. Indeed, if the isotropic, single-mass King models are adopted a priori as complete descriptions of the internal structures of Galactic globular clusters, then only *four* of these eleven variables are truly independent. These are (1) the mass-to-light ratio $\Upsilon_{V,0}$; (2) the concentration parameter $c = \log(r_t/r_0)$, which is directly related to the depth of a cluster’s potential well; and two other parameters—chosen here to be (3) binding energy and (4) total luminosity—required to normalize a dimensionless model to an observed object. The reduction to this physical basis requires seven definitions and model relations. (See also Djorgovski 1993a for an outline of this procedure, and King 1966 or Binney & Tremaine 1987 for more detailed descriptions of the models themselves.)

First, recall that, where they are available, the directly observed central velocity dispersions of Milky Way globulars (Pryor & Meylan 1993) have *already* been converted to King-model scale velocities σ_0 (in §2 above; see Fig. 1 and eq. [B1]). Also, the so-called core radii tabulated by Harris (1996; also Trager et al. 1993 and Djorgovski 1993a) are in fact the r_0 referred to here as scale radii (r_0 differs from the true projected half-power radius in low-concentration clusters).

Given these things, the definition of r_0 (King 1966) yields a cluster’s central *mass* density:

$$\Upsilon_{V,0} j_0 = 9\sigma_0^2 / (4\pi G r_0^2) , \quad (\text{A1})$$

where the basic scaling $\rho_0 \propto (\sigma_0^2 / G r_0^2)$ follows simply from dimensional analysis, but the coefficient $(9/4\pi)$ is specific to the King model. A central *luminosity* volume density alone is derived from an observed r_0 , c , and central surface brightness, by way of the trivial (model-independent) definition

$$\mu_{V,0} = 26.362 - 2.5 \log(I_0 / L_\odot \text{ pc}^{-2}) \quad (\text{A2})$$

and the model relation

$$I_0 = \mathcal{I}_0(c) j_0 r_0 , \quad (\text{A3})$$

where $\mathcal{I}_0$ is the dimensionless function of concentration shown in the bottom left panel of Fig. 1 and approximated by equation (B3). Equation (A1) is used in §3.1 to infer the central mass-to-light ratio $\Upsilon_{V,0}$ for the 39 non-core-collapsed clusters catalogued by Pryor & Meylan (1993).

The projected half-light radius, King scale radius, and central concentration are linked through the function

$$\mathcal{R}(c) = R_h/r_0 , \quad (\text{A4})$$

which is drawn in the bottom right panel of Fig. 1 and approximated in equation (B4) below. The average surface brightness within R_h is then given (for L in $L_\odot$ and R_h in pc) by

$$\langle \mu_V \rangle_h = 26.362 - 2.5 \log (L/2\pi R_h^2) , \quad (\text{A5})$$

where the total cluster luminosity is given in terms of c , j_0 and r_0 by

$$L = \mathcal{L}(c) j_0 r_0^3 . \quad (\text{A6})$$

(See Fig. 2 and eq. [B5] for the dimensionless function $\mathcal{L}$.) Finally, the cluster binding energy is expressed as (cf. eq. [2])

$$E_b = \frac{\sigma_0^4 r_0}{G} \mathcal{E}(c) = \left(\frac{9}{4\pi} \right)^{1/2} \frac{\sigma_0^5 (\Upsilon_{V,0} j_0)^{-1/2}}{G^{3/2}} \mathcal{E}(c) = \left(\frac{4\pi}{9} \right)^2 G (\Upsilon_{V,0} j_0)^2 r_0^5 \mathcal{E}(c) , \quad (\text{A7})$$

with $\mathcal{E}$ the function in equation (B2) and the top right panel of Fig. 1. (The last two equalities in eq. [A7] follow from the definition [A1].)

To repeat, then, a King-model globular cluster is defined fully by its core mass-to-light ratio, central concentration, binding energy, and total luminosity. Real clusters therefore can be expected to inhabit a nominally four-dimensional parameter space. However, the dimensionality of this space is reduced by one for every independent constraint on the four fundamental quantities. Thus, given relations of the type inferred in §3 for the Milky Way clusters,

$$\Upsilon_{V,0} = \text{constant} \quad \text{and} \quad E_b = A (L/L_\odot)^\gamma , \quad (\text{A8})$$

globulars constitute an essentially two-parameter family of objects (although, in principle, $\Upsilon_{V,0}$ or the $E_b(L)$ relation may further depend on external, environmental variables; in our Galaxy, $A \propto r_{\text{gc}}^{-0.4}$ according to §3.3 above.) That is, having specified L and c for an object (at some given r_{gc}), its binding energy E_b and mass-to-light ratio $\Upsilon_{V,0}$ follow immediately from equation (A8). The product $(j_0 r_0^3)$ is then given uniquely by equation (A6), and the combination $(j_0^2 r_0^5)$ by equation (A7). This sets the values of j_0 and r_0 individually, and all other cluster parameters follow from equations (A1)–(A5). If there exists a third basic relation, $c = c(\log L)$ —a situation which is strongly suggested, although not proven, by current data on the Galactic cluster system—then luminosity or mass alone (again, at a given r_{gc} in our Galaxy) controls the globular cluster sequence.

It is clear, as a result of this, that *any correlations between any observed or derived cluster parameters must trace back to some combination of the equations (A1)–(A7) characterizing King models, and/or to relations between the fundamental physical variables $\Upsilon_{V,0}$, c , E_b , and L .* Several examples of such observable correlations are now derived quite generally; their application to the Milky Way globular cluster system specifically is discussed in §§4 and 5 above.

A.1. Bivariate Correlations

Equation (A1) can be viewed either as a definition of r_0 or as an expression for $\Upsilon_{V,0}$ in terms of σ_0 , r_0 , and j_0 . Either way, it may be re-written as

$$(4\pi G/9) \Upsilon_{V,0} j_0 r_0 = \sigma_0^2 r_0^{-1} ;$$

or, in appropriate units,

$$\left(\frac{\sigma_0}{\text{km s}^{-1}} \right)^2 \left(\frac{r_0}{\text{pc}} \right)^{-1} = 5.9870 \times 10^{-3} \left(\frac{\Upsilon_{V,0}}{M_\odot L_\odot^{-1}} \right) \left(\frac{j_0 r_0}{L_\odot \text{pc}^{-2}} \right) .$$

Taking the logarithm of both sides and using equations (A2) and (A3) to introduce the observable $\mu_{V,0}$ then yields

$$\log \sigma_0 - 0.5 \log r_0 = -0.2 \mu_{V,0} - 0.5 \log \mathcal{I}_0(c) + 0.5 \log \Upsilon_{V,0} + 4.1610 . \quad (\text{A9})$$

Given a constant $\log \Upsilon_{V,0}$ (eq. [4]) and a very slowly varying $\mathcal{I}_0(c)$, this becomes an essentially bivariate correlation corresponding to one of Djorgovski's (1995) two equations for the fundamental plane of Galactic globular clusters (§4.2). It may also be put in terms of half-light cluster radii and average surface brightness by noting a relation between $\mu_{V,0}$ and $\langle \mu_V \rangle_h$ that follows from King-model definitions: Equations (A2), (A3), (A6), (A4), and (A5) combined give

$$\begin{aligned} \mu_{V,0} &= 26.362 - 2.5 \log \left[\frac{\mathcal{I}_0(c) j_0 r_0}{L_\odot \text{pc}^{-2}} \right] = 26.362 - 2.5 \log \left[2\pi \frac{\mathcal{I}_0(c) \mathcal{R}(c)^2}{\mathcal{L}(c)} \frac{L/2\pi R_h^2}{L_\odot \text{pc}^{-2}} \right] \\ &= \langle \mu_V \rangle_h - 2.5 \log (2\pi) - 2.5 \log \left[\frac{\mathcal{I}_0(c) \mathcal{R}(c)^2}{\mathcal{L}(c)} \right] , \end{aligned} \quad (\text{A10})$$

and substitution of this in equation (A9) yields

$$\log \sigma_0 - 0.5 \log R_h = -0.2 \langle \mu_V \rangle_h - 0.5 \log [\mathcal{L}(c)/\mathcal{R}(c)] + 0.5 \log \Upsilon_{V,0} + 4.5601 . \quad (\text{A11})$$

as another expression of equation (A1).

A power-law scaling of binding energy with total luminosity leads to another pair of equivalent bivariate correlations. With $E_b = A (L/L_\odot)^\gamma$ as in equation (A8), the first equality in equation (A7) reads

$$\sigma_0^4 r_0 = GA (L/L_\odot)^\gamma \mathcal{E}(c)^{-1} ,$$

where, again, the normalization A in general may vary with Galactocentric position (or with any other environmental factor appropriate to a given dataset). Applying equation (A6) and inserting the usual units, this becomes

$$\left(\frac{\sigma_0}{\text{km s}^{-1}} \right)^4 \left(\frac{r_0}{\text{pc}} \right)^{1-2\gamma} = 2.1558 \times 10^{-46} \left(\frac{A}{\text{erg}} \right) \left[\frac{\mathcal{L}(c)^\gamma}{\mathcal{E}(c)} \right] \left(\frac{j_0 r_0}{L_\odot \text{pc}^{-2}} \right)^\gamma ,$$

so that taking the logarithm and using the definition of $\mu_{V,0}$ (eqs. [A2] and [A3]) yields

$$\begin{aligned} \log \sigma_0 - 0.25 (2\gamma - 1) \log r_0 &= -(0.1 \gamma) \mu_{V,0} + 0.25 \log [\mathcal{L}(c)^\gamma / \mathcal{E}(c) \mathcal{I}_0(c)^\gamma] \\ &\quad + 0.25 \log A + 2.6362 \gamma - 11.4166 . \end{aligned} \quad (\text{A12})$$

And finally, a version that refers to half-light cluster quantities is obtained by applying equations (A4) and (A10):

$$\begin{aligned} \log \sigma_0 - 0.25 (2\gamma - 1) \log R_h &= -(0.1 \gamma) \langle \mu_V \rangle_h - 0.25 \log [\mathcal{E}(c) / \mathcal{R}(c)] \\ &\quad + 0.25 \log A + 2.8357 \gamma - 11.4166 , \end{aligned} \quad (\text{A13})$$

which is shown in §4.2 to correspond to Djorgovski’s (1995) second equation for the globular cluster fundamental plane in the Milky Way.

A.2. “Monovariate” Correlations

Given the suite of definitions listed above (eqs. [A1]–[A7]), plus the two fundamental-plane constraints of equation (A8), any globular cluster observables can be expressed as functions of central concentration and total luminosity (and any environmental factors, such as r_{gc} , which in general can be subsumed in eq. [A8]). Some of these relations can appear as monovariate cluster correlations if non-homology and variations in c are ignored.

One such expression follows from the third equality in equation (A7). Together with equations (A8) and (A6), this reads

$$\frac{E_b}{\text{erg}} = 1.6627 \times 10^{41} \left(\frac{\Upsilon_{V,0}}{M_\odot L_\odot^{-1}} \right)^2 \left(\frac{j_0}{L_\odot \text{pc}^{-3}} \right)^2 \left(\frac{r_0}{\text{pc}} \right)^5 \mathcal{E}(c) = \frac{A}{\text{erg}} \left(\frac{L}{L_\odot} \right)^{\gamma-2} \left[\frac{\mathcal{L}(c) j_0 r_0^3}{L_\odot} \right]^2 ,$$

or, after taking the logarithm of both sides,

$$\log r_0 = (2 - \gamma) \log L - \log [\mathcal{L}(c)^2 / \mathcal{E}(c)] - \log A + 2 \log \Upsilon_{V,0} + 41.2208 . \quad (\text{A14})$$

As §5 discusses, this does translate to a rough correlation between r_0 and L in the Milky Way (see Figs. 7 and 17). It also can be used to derive the dependence of $R_h = \mathcal{R}(c) r_0$ on L and c :

$$\log R_h = (2 - \gamma) \log L - \log [\mathcal{L}(c)^2 / \mathcal{E}(c) \mathcal{R}(c)] - \log A + 2 \log \Upsilon_{V,0} + 41.2208 , \quad (\text{A15})$$

accounting for the behavior of observed cluster half-light radii in Fig. 8 above. (Like r_0 , R_h is measured in pc.)

Correlations involving surface brightness follow from the basic definitions (A2), (A3), and (A6), which state that

$$\begin{aligned} \log r_0 &= 0.2 \mu_{V,0} + 0.5 \log (j_0 r_0^3) + 0.5 \log \mathcal{I}_0(c) - 5.2724 \\ &= 0.2 \mu_{V,0} + 0.5 \log L - 0.5 \log [\mathcal{L}(c) / \mathcal{I}_0(c)] - 5.2724 \end{aligned}$$

for j_0 in units of $L_\odot \text{ pc}^{-3}$ and r_0 in pc. This is itself a bivariate correlation; it is discussed further in §5, in connection with a rough correlation between r_0 and $\mu_{V,0}$ observed in the Milky Way cluster system (see Fig. 19). But it and equation (A14) also lead to

$$\log j_0 = (3\gamma - 5) \log L + \log \left[\mathcal{L}(c)^5 / \mathcal{E}(c)^3 \right] + 3 \log A - 6 \log \Upsilon_{V,0} - 123.662 \quad (\text{A16})$$

or to

$$\mu_{V,0} = 2.5 (3 - 2\gamma) \log L - 2.5 \log \left[\mathcal{L}(c)^3 \mathcal{I}_0(c) / \mathcal{E}(c)^2 \right] - 5 \log A + 10 \log \Upsilon_{V,0} + 232.466, \quad (\text{A17})$$

which corresponds to the correlation between $\log j_0 r_0$ and $\log L$ in Fig. 7 above. In addition, equations (A17) and (A10) together, or equations (A15) and (A5) together, imply

$$\langle \mu_V \rangle_h = 2.5 (3 - 2\gamma) \log L - 5 \log \left[\mathcal{L}(c)^2 / \mathcal{E}(c) \mathcal{R}(c) \right] - 5 \log A + 10 \log \Upsilon_{V,0} + 234.461. \quad (\text{A18})$$

Finally, a correlation between core velocity dispersion and total cluster luminosity can be obtained directly from the definition $E_b = \sigma_0^4 r_0 \mathcal{E}(c) / G$, the empirical scaling $E_b / \text{erg} = A (L / L_\odot)^\gamma$, and equation (A14):

$$\log \sigma_0 = 0.5(\gamma - 1) \log L - 0.5 \log [\mathcal{E}(c) / \mathcal{L}(c)] + 0.5 \log A - 0.5 \log \Upsilon_{V,0} - 21.7218 \quad (\text{A19})$$

for σ_0 in units of km s^{-1} . This version of the $E_b(L)$ relation in the Milky Way is compared to data in Fig. 18 above.

The six quantities on the left-hand sides of equations (A14)–(A19) are all expressed in terms of the King-model basis L , c , $\Upsilon_{V,0}$, and $E_b = A (L / L_\odot)^\gamma$ (once more, any effects of, say, Galactocentric position and metallicity are easily allowed to enter through these variables; see §5). Moreover, any correlation between any other set of globular cluster observables may be derived from equations (A14)–(A19)—the results of §A.1, for example, could be thus obtained. As for *any* collection of single-mass, isotropic King-model clusters, there is no *new* physical content in empirical scalings beyond those existing just among L , c , $\Upsilon_{V,0}$ and E_b . This follows necessarily from the fact that this chosen basis is (by definition) complete.

B. DIMENSIONLESS FUNCTIONS OF CENTRAL CONCENTRATION

This Appendix lists ad hoc polynomial fits that describe the variation of certain derived quantities in single-mass, isotropic King (1966) models, as functions of the central concentration $c \equiv \log(r_t / r_0)$. These fits have been obtained by comparing with numerical integrations of King models with $0.12 \lesssim c \lesssim 3.6$, or $0.5 \leq W_0 \leq 16$ for W_0 the (normalized) central potential. The fits generally will *not* be reliable if extrapolated beyond this range in c , but all Galactic globular clusters are described by $0.5 \leq c \leq 2.5$.

The projected, or line-of-sight, velocity dispersion at the center of a model cluster is given in terms of the velocity *scale parameter* σ_0 —which differs from the true one-dimensional velocity dispersion for low-concentration models—by

$$\log (\sigma_{p,0}/\sigma_0) = -14.203 \log \left[1 + 0.11313 \times 10^{-1.1307 c} \right] , \quad (\text{B1})$$

which deviates from the curve in Fig. 1 above by less than 0.005 dex for $c \gtrsim 0.5$.

The dimensionless binding energy (eq. [2] above, and the top right panel of Fig. 1) may be fit by

$$\begin{aligned} \log \mathcal{E} \equiv \log \left(GE_b/\sigma_0^4 r_0 \right) = & -1.64893 + 2.82056 c + 9.38926 c^2 - 26.0275 c^3 + 30.6474 c^4 \\ & -20.7951 c^5 + 8.61423 c^6 - 2.13978 c^7 + 0.291982 c^8 \\ & -0.0167994 c^9 , \end{aligned} \quad (\text{B2})$$

with an absolute error of $\lesssim 0.01$ dex (relative error $\lesssim 1\%$) for $c \gtrsim 0.5$.

The central intensity, or surface density, in terms of the central volume density j_0 and the King radius r_0 , is (cf. Djorgovski 1993a, and the bottom left panel of Fig. 1 above)

$$\log \mathcal{I}_0 \equiv \log (I_0/j_0 r_0) = 0.3022 - 7.5726 \log \left[1 + 0.2180 \times 10^{-1.1291 c} \right] , \quad (\text{B3})$$

to within an absolute error of $\lesssim 0.005$ dex in the interval $0.5 \lesssim c \lesssim 3.6$. [Note that in the limit $c \rightarrow \infty$, this formula gives $\mathcal{I}_0 \rightarrow 2.005$, slightly lower than the correct value for an isothermal sphere, $\mathcal{I}_0(\infty) = 2.018$.] For these single-mass models, which have spatially constant mass-to-light ratios, $\mathcal{I}_0$ is also equal to the dimensionless central *mass* surface density, $\Sigma_0/\rho_0 r_0$.

The *projected* half-light radius, R_h , is related to the King radius, $r_0 \equiv (9\sigma_0^2/4\pi G\rho_0)^{1/2}$, by

$$\begin{aligned} \log \mathcal{R} \equiv \log (R_h/r_0) = & -0.602395 + 1.36023 c - 1.67086 c^2 + 2.65848 c^3 - 2.71152 c^4 \\ & +1.42555 c^5 - 0.274551 c^6 - 0.0381277 c^7 + 0.0217849 c^8 \\ & -0.00225252 c^9 , \end{aligned} \quad (\text{B4})$$

which differs by no more than 0.007 dex from the $\log \mathcal{R}$ obtained by numerically integrating any King model with $0.12 \lesssim c \lesssim 3.6$ (bottom right of Fig. 1 above).

Finally, the dimensionless total luminosity (top panel of Fig. 2) is given by

$$\begin{aligned} \log \mathcal{L} \equiv \log \left(L/j_0 r_0^3 \right) = & -0.725444 + 1.90743 c + 4.63720 c^2 - 13.0266 c^3 + 14.8724 c^4 \\ & -9.43699 c^5 + 3.39098 c^6 - 0.578169 c^7 - 0.00512814 c^8 \\ & +0.0162534 c^9 - 0.00165078 c^{10} . \end{aligned} \quad (\text{B5})$$

The relative error of this expression is $\lesssim 0.5\%$, in $\log \mathcal{L}$, for any $0.5 \lesssim c \lesssim 3.6$; it may be compared to the fit given by Djorgovski (1993a), which applies over a more restricted range in central concentration. As with equation (B3), $\mathcal{L}$ also gives the dimensionless *mass*, $M/\rho_0 r_0^3$, of a single-mass cluster.

REFERENCES

- Bellazzini, M. 1998, *New Astronomy*, 3, 219
- Bellazzini, M., Vesperini, E., Ferraro, F. R., & Fusi Pecci, F. 1996, *MNRAS*, 279, 337
- Bender, R., Burstein, D., & Faber, S. 1992, *ApJ*, 399, 462
- Binney, J., & Tremaine, S. 1987, *Galactic Dynamics* (Princeton: Princeton University Press)
- Brosche, P., & Lentes, F.-T. 1984, *A&A*, 139, 474
- Burstein, D., Bender, R., Faber, S., & Nolthenius, R. 1997, *AJ*, 114, 1365
- Busarello, G., Capaccioli, M., Capozziello, S., Longo, G., & Puddu, E. 1997, *A&A*, 320, 415
- Djorgovski, S. 1993a, in *ASP Conf. Ser. 50, Structure and Dynamics of Globular Clusters*, ed. S. Djorgovski and G. Meylan (San Francisco: ASP), 373
- Djorgovski, S. 1993b, in *ASP Conf. Ser. 48, The Globular Cluster–Galaxy Connection*, ed. G. H. Smith and J. P. Brodie (San Francisco: ASP), 496
- Djorgovski, S. 1995, *ApJ*, 438, L29
- Djorgovski, S., & Davis, M. 1987, *ApJ*, 313, 59
- Djorgovski, S., & Meylan, G. 1994, *AJ*, 108, 1292
- Dressler, A., Lynden-Bell, D., Burstein, D., Davies, R. L., Faber, S. M., Terlevich, R. J., & Wegner, G. 1987, *ApJ*, 313, 42
- Graham, A., & Colless, M. 1997, *MNRAS*, 287, 221
- Gunn, J. E., & Griffin, R. F. 1979, *AJ*, 84, 752
- Harris, W. E. 1996, *AJ*, 112, 1487
- Harris, W. E., & Pudritz, R. E. 1994, *ApJ*, 429, 177
- King, I. R. 1966, *AJ*, 71, 64
- Kormendy, J. 1985, *ApJ*, 295, 73
- Kritsuk, A. G. 1997, *MNRAS*, 284, 327
- Mandushev, G., Spassova, N., & Staneva, A. 1991, *A&A*, 252, 94
- McLaughlin, D. E., & Pudritz, R. E. 1996, *ApJ*, 457, 578
- Michie, R. W. 1963, *MNRAS*, 125, 127
- Murray, S. D., & Lin, D. N. C. 1992, *ApJ*, 400, 265
- Ostriker, J. P., & Gnedin, O. Y. 1997, *ApJ*, 487, 667
- Pahre, M. A., de Carvalho, R. R., & Djorgovski, S. G. 1998, *AJ*, 116, 1606
- Pryor, C., & Meylan, G. 1993, in *ASP Conf. Ser. 50, Structure and Dynamics of Globular Clusters*, ed. S. G. Djorgovski and G. Meylan (San Francisco: ASP), 357

- Richstone, D. O., & Tremaine, S. 1986, *AJ*, 92, 72
- Rosenberg, A., Saviane, I., Piotto, G., & Held, E. V. 1998, *A&A*, 339, 61
- Saito, M. 1979, *PASJ*, 31, 181
- Schaeffer, R., Maurogordato, S., Cappi, A., & Bernardeau, F. 1993, *MNRAS*, 263, L21
- Shapley, H., & Sawyer, H. B. 1927, *Harvard College Obs. Bull.*, No. 849
- Trager, S. C., Djorgovski, S., & King, I. R., 1993, in *ASP Conf. Ser. 50, Structure and Dynamics of Globular Clusters*, ed. S. Djorgovski and G. Meylan (San Francisco: ASP), 347
- Trager, S. C., King, I. R., & Djorgovski, S. 1995, *AJ*, 109, 218
- van den Bergh, S. 1994, *ApJ*, 435, 203
- van den Bergh, S. 1995, *AJ*, 110, 1171
- van den Bergh, S., Morbey, C., & Pazder, J. 1991, *ApJ*, 375, 594
- van der Marel, R. P. 1991, *MNRAS*, 253, 710
- Vesperini, E. 1997, *MNRAS*, 287, 915

Fig. 1.— Dimensionless functions of central concentration for single-mass, isotropic King (1966) models. Analytic expressions for each are given in Appendix B. The observed concentrations of Milky Way globulars lie in the range $0.5 \lesssim c \lesssim 2.5$.

Fig. 2.— *Top panel:* Dimensionless total luminosity $\mathcal{L}$ of King models, as a function of central concentration. See also equation (B5). *Bottom panel:* Comparison of model luminosities $\mathcal{L}(c)j_0r_0^3$ to the directly observed $L = 0.4(4.83 - M_V)$ for globulars from Harris (1996). Filled circles are regular, King-model clusters; open squares are post-core-collapse objects. Straight line is the equality $\log L_{\text{mod}} = \log L$, and the typical observational errorbar on $\log L_{\text{mod}}$ is shown in the upper left corner.

Fig. 3.— Determination, according to equation (3), of the mean core mass-to-light ratio of globular clusters in the Milky Way. Filled circles correspond to 39 King-model clusters tabulated by Pryor & Meylan (1993); open circles have ρ_0 computed for these objects using the directly observed $\sigma_{p,0}$ instead of the model velocity scale σ_0 . Open squares represent PCC clusters. The solid line has the equation $\log \rho_0 = 0.16 + \log j_0$, obtained from a least-squares fit of the regular-cluster data only. The errorbars at upper left show typical observational uncertainties in ρ_0 and j_0 .

Fig. 4.— Concentration parameters and binding energies, as functions of total luminosity, for globulars with measured central velocity dispersions (Pryor & Meylan 1993). Filled circles correspond to regular, King-model clusters; open squares, to PCC objects. The three lower rows present the results of calculating E_b according to each of equations (5). Integrated absolute magnitudes are taken from the catalogue of Harris (1996), while total masses are obtained by Pryor & Meylan (1993) from fits of multi-mass, anisotropic King models.

Fig. 5.— Confirmation that the binding energy computed using equation (5c) (with $\log \Upsilon_{V,0} \equiv 0.16$) is equal to that obtained from (5a) for the regular globulars (filled circles) in the catalogue of Pryor & Meylan (1993). Open squares again correspond to the PCC clusters in their sample. This plot is equivalent to Fig. 3.

Fig. 6.— Central concentrations and binding energies of 109 King-model globulars and 30 PCC clusters (filled circles and open squares) in the catalogue of Harris (1996). The straight line in the top panel traces the rough relation $c = -0.57 + 0.4 \log L$. The solid line in the bottom panel represents the least-squares fit $\log E_b = 39.89 + 2.05 \log L$; dashed lines show the 3σ limits (from eq. [7]), $\log E_b = 41.09 + 1.8 \log L$ and $\log E_b = 38.69 + 2.3 \log L$.

Fig. 7.— Monovariate correlations $\log r_0$ vs. $\log L$ and $\log j_0 r_0$ vs. $\log L$ for Galactic globulars in the Harris (1996) catalogue, as poor reflections of the more fundamental correlation between $\log E_b$ and $\log L$. Filled circles are regular clusters; open squares are PCC objects. The straight line in the top panel has $\log r_0 = 1.63 - 0.3 \log L$; in the middle panel, $\log j_0 r_0 = -2.86 + 1.25 \log L$; in the bottom panel, $\log j_0^2 r_0^5 = -1.33 + 1.7 \log L$.

Fig. 8.— Dependence of cluster half-light radius on luminosity and Galactocentric radius. Point

types are the same as in Fig. 7. R_h is seen to increase systematically with r_{gc} (the line in the top right panel is $\log R_h = 0.226 + 0.4 \log r_{gc}$), but is essentially independent of L at a given position in the Milky Way. R_h^* is the normalized quantity $R_h(r_{gc}/8 \text{ kpc})^{-0.4}$; the dashed line in the bottom plot has $\log R_h^* \equiv 0.59$.

Fig. 9.— Insensitivity of three of the basic King-model cluster parameters to the “environmental” factors of Galactocentric position and cluster metallicity. Open squares again represent PCC clusters; note that c has been *arbitrarily* set to 2.5 for many of these.

Fig. 10.— Dependence of cluster binding energy on Galactocentric radius. Straight line in the bottom panel represents the fit $E_b L^{-2.05} \propto r_{gc}^{-0.4}$, as in equation (8), for regular clusters from the Harris (1996) catalogue (filled circles). The observational errorbar shown for $\log E_b$ is ± 0.5 dex.

Fig. 11.— Independence of binding energy and metallicity for the 139 clusters taken from Harris (1996).

Fig. 12.— Concentrations c and normalized binding energies $E_b^* \equiv E_b(r_{gc}/8 \text{ kpc})^{-0.4}$ for regular and PCC Galactic globulars within and beyond $r_{gc} = 8 \text{ kpc}$. Dashed lines in both top panels are $c = -1.03 + 0.5 \log L$ and $c = -2.44 + 0.8 \log L$. The r.m.s. scatter in c is $\Delta = 0.29$ dex (excluding Palomar 1) at $r_{gc} > 8 \text{ kpc}$, and $\Delta = 0.35$ for $r_{gc} < 8 \text{ kpc}$. Solid line in the bottom panels is the fit $\log E_b^* = 39.86 + 2.05 \log L$, and dashed lines there are the 3σ limits (from eq. [8]) $\log E_b^* = 41.06 + 1.8 \log L$ and $\log E_b^* = 38.66 + 2.3 \log L$. The r.m.s. scatter in $\log E_b^*$ is ± 0.5 dex.

Fig. 13.— Fundamental plane of Galactic globular clusters (109 regular and 30 core-collapsed) in the catalogue of Harris (1996). The orthogonal axes in ϵ -space are defined in equation (11). The (ϵ_1, ϵ_2) and (ϵ_1, ϵ_3) cross-sections of this volume provide edge-on views of the fundamental plane, defined by $\epsilon_1 = 39.86$ (eq. [8], drawn as bold solid lines). A face-on view of the fundamental plane is found in the (ϵ_2, ϵ_3) slice of ϵ -space. Clusters do not fill this plane uniformly but are confined, particularly at large galactocentric distances, to a fairly narrow band within it. The dashed line is a fit $\epsilon_3 = -12.5 + 0.13 \epsilon_2$ to the globulars (excluding Palomar 1) with $r_{gc} > 8 \text{ kpc}$. Most of the PCC clusters appear to fall away from the fundamental plane here, but these objects are generally *not* well described by King models.

Fig. 14.— *Left panels*: Effects of cluster non-homology on two empirical, bivariate correlations involving *half-light* parameters of Galactic globulars. The functions $\mathcal{L}(c)$, $\mathcal{R}(c)$, and $\mathcal{E}(c)$ are given individually in Figs. 1 and 2, and in Appendix B. Points are placed at the observed c -values of the 39 regular clusters in the catalogue of Pryor & Meylan (1993). *Right panels*: Observed correlations in the Pryor & Meylan cluster sample. Filled circles are their regular clusters, and the 17 open squares are their PCC clusters. Open circles plot the uncorrected $\sigma_{p,0}$ for the regular objects, rather than the model velocity scales σ_0 . Solid lines in the top and bottom panels represent equations (12b) and (13b); dashed lines are least-squares fits to the data (filled circles). The distant cluster NGC 2419 is noted as an outlier (see Fig. 18).

Fig. 15.— *Left panels:* Effects of cluster non-homology on two empirical, bivariate correlations involving *core* parameters of Galactic globulars. The functions $\mathcal{I}_0(c)$, $\mathcal{L}(c)$, and $\mathcal{E}(c)$ are given individually in Figs. 1 and 2, and in Appendix B. The dashed line in the bottom panel shows the rough scaling $0.25 \log (\mathcal{L}^{2.05}/\mathcal{E}\mathcal{I}_0^{2.05}) \approx \text{const.} + 0.17c$. *Right panels:* Correlations observed for the Pryor & Meylan (1993) clusters. Point types have the same meaning as in Fig. 14. Solid line in the top panel represents equation (12a) and is indistinguishable from a least-squares fit to the regular-cluster data. Solid line in the bottom panel is equation (13a) with an intercept of 4.147, estimated by assuming a constant $\langle 0.25 \log (\mathcal{L}^{2.05}/\mathcal{E}\mathcal{I}_0^{2.05}) \rangle = 0.204$ for all 39 King-model clusters. The least-squares fit (dashed line) is quite different ($\log \sigma_0 - 0.775 \log r_0 = -0.246 \mu_{V,0} + 4.860$) because of the significant non-homology in this case.

Fig. 16.— Virial ratio $2E_k/E_b$ as a function of radius inside three single-mass, isotropic King (1966) models bracketing the range of central concentration observed for Galactic globulars. Cluster cores, defined by $r \leq r_0$, in general do *not* obey a simplified virial theorem of the form $2E_k = E_b$.

Fig. 17.— Source of the monovariate correlation between $\log r_0$ and $\log L$ in Fig. 7. *Top panel:* The non-homology term $\log (\mathcal{L}^2/\mathcal{E})$ as a function of central concentration. Solid curve is obtained from numerical integrations of King models, points are placed at the observed c -values of the 109 regular clusters in the catalogue of Harris (1996), and the dashed line shows $\log (\mathcal{L}^2/\mathcal{E}) \approx \text{const.} + 0.7c$. *Bottom panel:* The full correlation involving $\log r_0$, c , $\log L$, and $\log r_{\text{gc}}$. Data for 30 PCC clusters from Harris (1996) are also shown (open squares). Solid line is equation (14a): $\log r_0 + \log (\mathcal{L}^2/\mathcal{E}) = 1.681 - 0.05 [\log L - 8 \log (r_{\text{gc}}/8 \text{ kpc})]$; the dashed line, with an intercept of 1.63, is the least-squares fit to the regular-cluster data (filled circles).

Fig. 18.— *Top panel:* Monovariate correlation between $\log \sigma_0$ and $\log L$ for clusters from Pryor & Meylan (1993). Point types have the same meaning as in Fig. 14. The observational errorbar on $\log \sigma_0$ is ± 0.09 dex. The straight line is equation (14f) after substituting appropriate averages for the ratio $\mathcal{E}(c)/\mathcal{L}(c)$ and r_{gc} (see text). *Bottom panel:* The full correlation expected on the basis of the fundamental-plane constraints on $\Upsilon_{V,0}$ and E_b^* . Solid line is equation (14f); dashed line is a least-squares fit to the regular-cluster data (filled circles).

Fig. 19.— *Top left:* Monovariate correlation between $\log r_0$ and $\mu_{V,0}$ for globulars in the catalogue of Harris (1996). Solid points refer to the 109 King-model clusters; open squares, to the 30 PCC objects. The observed r.m.s. scatter in $\log r_0$ exceeds the observational errorbar of ± 0.1 dex. *Top right:* Real variations in the “core mass,” $M_0 \propto j_0 r_0^3$. *Bottom left:* Deviations from the line expected if core mass were a constant are not random, but correlated with total luminosity. *Bottom right:* The first three panels are explained as results of the behavior of *total* luminosity as a function of c in King models, combined with the definition of core surface brightness. The straight line is equation (15).




Fig. 1.—




Fig. 2.—




Fig. 3.—




Fig. 4.—




Fig. 5.—




Fig. 6.—




Fig. 7.—




Fig. 8.—




Fig. 9.—




Fig. 10.—




Fig. 11.—




Fig. 12.—




Fig. 13.—




Fig. 14.—




Fig. 15.—




Fig. 16.—




Fig. 17.—




Fig. 18.—




Fig. 19.—